



Product Name: APPARENT MCPA 750 SELECTIVE HERBICIDE
APVMA Approval No: 68355/123018



| | |
|-------------------------|---|
| Label Name: | APPARENT MCPA 750 SELECTIVE HERBICIDE |
| Signal Headings: | POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING |
| Constituent Statements: | ACTIVE CONSTITUENT: 750 g/L MCPA present as the dimethylamine salt SOLVENT: 280 g/L DIMETHYLACETAMIDE |
| Mode of Action: | GROUP I HERBICIDE |
| Statement of Claims: | For the control of broadleaf weeds in cereals, linseed, poppies, pastures, rice, sugar cane and turf as specified in the Directions for Use table |
| Net Contents: | 5L-1000L |
| Restrains: | Restrains: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions. DO NOT apply if rain is likely within 6 hours. |
| Directions for Use: | This section contains file attachment. |
| Other Limitations: | |
| Withholding Periods: | WITHHOLDING PERIOD: Cereal grain: DO NOT harvest, graze or cut for stock food for 7 days after application. Pasture: DO NOT graze or cut for stock food for 7 days after application. Poppies: DO NOT harvest for 5 weeks after application. |

| | |
|-----------------------|---|
| | Harvest Withholding Period (other than poppies and cereal grain): NOT required when used as directed. |
| Trade Advice: | |
| General Instructions: | This section contains file attachment. |
| Resistance Warning: | <p>RESISTANT WEEDS WARNING GROUP I HERBICIDE APPARENT MCPA 750 Selective Herbicide is a member of the Phenoxy group of herbicides. This product has the disruptors of plant cell growth mode of action. For weed resistance management this product is a Group I Herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR APPARENT Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.</p> |
| Precautions: | |
| Protections: | <p>PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto healthy susceptible plants/crops, cropping lands or pastures. Drift Warning: DO NOT use on or in situations where damage to susceptible crops – such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, Belahs, Eucalypts) – may occur from direct application, spray drift or vapours. LEGUME TOLERANCE The use of MCPA is preferred to 2,4-D Amine (Amicide) or Ester (Estercide) in situations where legumes are undersown in the crop. Note: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate dams, rivers or streams with the product or used containers.</p> |
| Storage and Disposal: | <p>STORAGE AND DISPOSAL</p> <p>Non-recyclable Containers: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.</p> |

| | |
|--|---|
| | <p>Envirodrum Containers: Store in the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any water or any other foreign matter. After each use of the product, please ensure the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drums remain the property of AIRR APPARENT Pty Ltd.</p> <p>Refillable Containers: Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs AIRR APPARENT Pty Ltd should be notified immediately. This minibulk/bulk container is reusable and remains the property of AIRR APPARENT Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return AIRR APPARENT Pty Ltd for cleaning, relabelling and refilling.</p> |
|--|---|

| | |
|---------------------------|---|
| Safety Directions: | <p>SAFETY DIRECTIONS Harmful if swallowed. Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product is on skin, immediately wash area with soap and water. If product or spray is in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.</p> |
|---------------------------|---|

| | |
|--------------------------------|---|
| First Aid Instructions: | <p>FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p> |
|--------------------------------|---|

| | |
|----------------------------|--|
| First Aid Warnings: | |
|----------------------------|--|

DIRECTIONS FOR USE

1. FIELD CROPS

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|--------------------------------------|--|--------------------|---|--|
| Wheat, Barley, Oats, Triticale | Refer to Weed Table | TAS, SA only | 460mL-1.45L | Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing. |
| | | QLD only | | Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem). |
| | | NSW, ACT only | | Apply 460mL rate after crop has reached 5 leaf stage and weeds marked # in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. |
| | | VIC only | 460-960mL | Apply 460mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered. |
| | | WA only | 600mL-1.35L | Apply to crop from 5 leaf to flag leaf emergence (Zadoks 15-37). |
| Wheat, Barley, Triticale, Cereal Rye | Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish | WA only | 270mL plus 350mL diuron (500g/L flowable) | CROP: 3-4 leaf stage (Zadoks 13-14). |
| | | | 330mL plus 500mL diuron (500g/L flowable) | CROP: 4-5 leaf stage (Zadoks 14-15). Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics. |
| | Amsinckia (Yellow Burr), Capeweed, Charlock Deadnettle, Double Gee (Spiny Emex, Three Corner Jack), <i>Erodium</i> sp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip | NSW, ACT, VIC only | 330mL plus 500mL diuron (500g/L flowable) | Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW, ACT and SA only) and 5 leaf stage (VIC only). Weeds must be in the 2-5 leaf stage. DO NOT apply to undersown medic or lucerne. DO NOT use on Kamilaroi, Kite, Olympic or Shortim Wheat. |
| | Capeweed, Mustard, Three Corner Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel | SA only | 230mL plus 500mL diuron (500g/L flowable) | DO NOT use on very light sandy soils. |
| | Soldier Thistle | SA only | 460mL plus 200mL Apparent Clopyralid 300 | Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing. |
| Skeleton Weed | 660mL plus 500mL Apparent Clopyralid 300 | | Apply to rosettes from 5-15cm diameter when the crop is between the leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August. This rate will provide control of Skeleton weed until harvest and will substantially reduce seed numbers the following season. | |

| | | | | |
|------------|--|------------------------------|--|---|
| Field Peas | Prickly Lettuce (<i>Lactuca serriola</i>) | NSW, VIC, TAS, SA only | <u>Weed Stage:</u> 4 leaf, 80mL plus 125mL diflufenican (500g/L) | FIELD PEAS: Apply early post-emergence after the third node stage and before the start of flowering. Warning: Field peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican (500g/L). APPLICATION AND WEED CONTROL: Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: <ul style="list-style-type: none"> • high crop and weed density, • late season germinations, • abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required. |
| | | | <u>Weed Stage:</u> 6 leaf, 100mL plus 150mL diflufenican (500g/L) | |
| | Charlock (Wild Mustard) (<i>Sinapis arvensis</i>) | NSW, VIC, TAS, SA only | <u>Weed Stage:</u> 4 leaf, 80mL plus 125mL diflufenican (500g/L) | |
| | | | <u>Weed Stage:</u> 6 leaf, 100mL plus 150mL diflufenican (500g/L) | |

1. FIELD CROPS – continued

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|---|---|-----------------------------|----------------|--|
| Linseed | Refer to Weed Table | QLD only | 460mL-1.45L | Apply when crop is 10-15cm tall with at least 170L/ha of water. DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft. |
| | | NSW, ACT, VIC, SA, TAS only | 460-730mL | |
| Oilseed poppies (<i>Papaver somniferum</i>) | Fat Hen (<i>Chenopodium album</i>), Spear Thistle (<i>Cirsium vulgare</i>) | NSW, TAS only | 1L/ha | Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30-40L of water/ha. This treatment will control late germinating weeds if they are green and at the bud or early flower stage. Spraying at this time will twist weeds below the crop canopy. DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops. |
| Sugar Cane | Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory | QLD only | 930mL | Post-emergent: Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties. |
| | As above plus Fat Hen, Noogoora Burr | | 1.45L | |
| Rice | Dirty Dora, Docks, Starfruit, Sedges, Buddah (Butter) Pea | ALL STATES | 485mL to 970mL | Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485mL/ha at 35 days after aerial sowing (approx. 2 tillers per rice plant). Use 970mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. Water Management: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying. |
| Flax | Refer to Weed Table | VIC, TAS only | 325mL | Apply to crop when 8-20cm high. |

2. PASTURES, SEED CROPS & TURF

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|----------------------------------|--|----------------------------|--|--|
| Grass Pastures | Refer to Weed Table | ALL STATES | 460mL-2.7L | Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume Tolerance" below. Note: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying. |
| | Salvation Jane | SA only | 660mL plus 400mL diuron (500g/L flowable) | Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm diameter. |
| | Soldier Thistle | | 460mL-1.45L | Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 575mL. |
| | Seedling and established Horehound | NSW, ACT only | 2.0L | Apply in autumn when horehound plants are at the seedling stage. If sub-clover or medic plants are present they will be damaged by this application. |
| Grass Seed Crops | Refer to Weed Table | QLD, VIC, TAS, SA, WA only | 460mL-2.7L | Apply to established grass seed crops. |
| Sub-Clover Seed Crops | Refer to Weed Table | VIC, SA only | 460-960mL | DO NOT exceed rate specified. Spray in autumn and early winter. |
| Pastures – Spray/Graze Technique | Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip | NSW, ACT, VIC only | 230-930mL | Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. See "Legume Tolerance" under General Instructions. Precaution: An increased quantity of poisonous plants may be eaten by stock using spray/graze e.g. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's Curse. Observe withholding period of at least 7 days after spraying. |
| | | TAS only | 460mL | |
| | Refer to Weed Table | WA only | 610mL-1.25L | |
| | Capeweed, Dandelion, Seedling Dock, Three Corner Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip | SA only | 330mL | |
| | Soldier Thistle | | 395mL | |
| Turf | Refer to Weed Table | ALL STATES | 330mL or 2.8mL per 100 sq. metres | Newly laid: Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses. |
| | | | 930mL - 1.8L or 10-19mL per 100 sq. metres | Established: Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses. |

3. SPOT SPRAYING

| SITUATION & CROP | WEEDS CONTROLLED | STATE | MIXING RATES/COMMENTS |
|-----------------------------|-------------------------|--------------|---|
| High Volume Spraying | Refer to Weed Table | ALL STATES | Use 130mL/100L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 270mL/100L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. |
| Knapsack Application | Refer to Weed Table | ALL STATES | Use 13mL/10L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 27mL/10L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. |

4. NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

| SITUATION & CROP | WEEDS CONTROLLED | STATE | RATE/ha | CRITICAL COMMENTS |
|---|-------------------------|--------------|---------------------|---|
| Non-Crop areas, Fallow Land, Industrial and Commercial Areas, Rights-of-way | Refer to Weed Table | ALL STATES | Refer to Weed Table | Apply at rates shown in Weed Table when weeds only are present and non-crop or pasture is involved. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture situation, use only the rate given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

| WEEDS CONTROLLED | APPLICATION RATE PER HECTARE | | | | | | CRITICAL COMMENTS |
|---------------------------|------------------------------|---------------|-------------|-------------|-------------|-------------|--|
| | VIC | NSW, ACT | SA | TAS | QLD | WA | |
| Ball Mustard | - | - | 1.15L | - | - | - | Apply at rosette stage. |
| Bathurst Burr | 1.35L | 660mL-1.35L | 1.35L | 1.35L | 1.35L | 1.35L | Spray young seedlings only. |
| Black Bindweed | - | 970mL-1.35L | - | - | 930mL-1.35L | - | Apply at young stage prior to flowering. |
| California Burr | - | 660mL-1.35L | - | 1.45L | - | - | Spray young seedlings only. |
| Cape Tulip | - | - | - | - | - | 1.35L | Apply just prior to flowering. Gives mild suppression only. |
| Capeweed | - | 1.45L | 1.45L | 1.45L | 1.45L | - | Apply at seedling stage. |
| Carrot Weed | 1.35L | - | - | - | 1.35L | - | Apply at seedling stage. |
| Charlock | 460mL | 660mL | 460mL | 660mL-1.45L | 460mL | - | Spray up to rosette stage. |
| Cutleaf Mignonette | - | - | 1.45L | - | - | - | Apply at young rosette stage. |
| Dandelion | 1.35L | 1.35L | 1.35L | 1.35L | 1.35L | 1.35L | Apply at young rosette stage. |
| Deadnettle | - | 1.45L | - | - | 1.45L | - | Moderately susceptible. Apply at seedling stage. |
| Docks | 960mL | - | 1.45L | 1.45L | - | - | Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected. |
| Fat Hen | 1.35L | 660mL-1.35L | - | 1.35L | 1.35L | - | Spray up to rosette stage. |
| Fennel | 1.80L | 1.80L | 1.80L | 1.80L | 1.80L | - | Apply at young stage. |
| Field Bindweed | 1.35L | - | 1.35L | 1.35L | 1.35L | 1.35L | Apply at bud stage, repeating over several seasons. |
| Field Cress | - | - | - | 1.45L | - | - | Apply at young stage. |
| Flatweed or Catsear | - | - | - | 1.45L | - | 930mL | Apply when weeds are 10-15cm high. |
| Fumitory (red) | 930mL | 930mL | 930mL | - | 930mL | - | Apply at young stage prior to flowering. |
| Hoary Cress | 3.0L | - | 1.45L | - | 3.0L | - | Apply before flowering. |
| Horehound | - | 2.0L | 1.45L | - | - | - | Apply at rosette stage. |
| Hedge Mustard | 730mL# | - | 460mL | 460mL | 460mL | 825mL | Apply at rosette stage. |
| Hexham Scent or Melilotus | 960mL | 800mL | 930mL | - | 930mL | - | Apply at young seedling stage. |
| Lesser Swinecress | - | - | - | 1.45L | - | - | Apply at young stage. |
| Lincoln Weed | - | - | 460mL | - | - | - | Apply at rosette stage. |
| London Rocket | - | - | - | - | - | 825mL | - |
| Lupins | - | 460mL - 960mL | - | - | - | 825mL | Spray up to 10cm high. |
| Mintweed | 1.35L | 1.35L | - | - | 1.35L | - | Spray young seedlings only. |
| Mustards | 730mL# | 660mL# | 730mL | 730mL | - | 825mL | Spray up to rosette stage. |
| Noogoora Burr | 1.35mL | 660mL-1.35L | 1.35L | - | 1.35L | 1.35L | Spray young seedlings only. |
| Opium Poppy | - | - | - | 1.45L | - | - | Apply at young stage. |
| Paterson's Curse | - | 660-960mL | - | - | 2.70L | - | Apply early rosette stage. |
| Pimpernel | - | - | 460mL | - | - | - | Apply at rosette stage. |
| Plantains | - | 1.35L | - | 1.35L | 1.35L | - | Apply at young stage. |
| Rapeseed | - | 660-960mL | - | - | - | - | Spray up to rosette stage. |
| Rough Poppy | - | 460-960mL | - | - | - | - | Spray up to rosette stage. |
| Safflower | - | 460-660mL | - | - | - | - | Spray up to rosette stage |
| Shepherd's Purse | - | - | - | 1.45L | - | - | Apply at rosette stage. |
| Skeleton Weed | - | 960mL-1.35L | - | - | - | - | Spray rosettes before aerial growth commences. |
| Sorrel | - | - | 1.45L | - | - | - | Apply to rosette stage. |
| Stinkwort | 1.35L | 660-960mL | 1.35L | - | - | 1.35L | Apply when weeds are 10-15cm high. |
| Sunflower (seedlings) | - | 690mL-1L | - | - | - | - | Spray up to rosette stage. |
| Thistles : | | | | | | | |
| - Artichoke | 1.05L | - | 1.45L | - | 1.35L | 460mL | Apply at rosette stage. |
| - Nodding | - | 1.65L | - | - | - | - | Spray early rosettes. Repeat spray required in following spring or autumn. |
| - Saffron | 960mL-1.65L | 660mL-1.35L | 930mL-1.65L | 930mL-1.65L | 930mL-1.65L | 960mL-1.65L | Spray up to rosette stage (September - WA). |
| Scotch or Cotton | 1.45L | - | 460-930mL | 1.45L | - | - | Apply at seedling stage. |
| Slender | 2.0L | 660mL-1.35L | 660mL | 2.0L | - | 2.0L | Spray young seedlings only. |
| Soldier | 730mL | - | 730mL | - | - | - | Spray young seedlings. Use higher rate for larger weeds. |
| Spear | - | 960mL-1.35L | - | 1.45L | - | - | Apply at rosette stage. |
| Star | - | - | 730-930mL | - | - | - | Apply at seeding stage. Use higher rate for larger weeds. |
| Variogated | 960mL | 660mL-1.35L | 460-930mL | 960mL | 960mL | 960mL | Spray young seedlings only. Use 1.35L/ha (SA) for well-developed weeds. |
| Turnip Weed | 730mL# | 660mL# | 660mL | - | - | - | Spray up to rosette stage. |
| Wild Artichoke | - | - | 1.45L | - | - | - | Apply at rosette stage. |
| Wild Radish | 930mL | 660mL# | 930mL | 1.35L | 460-730mL | 800mL | Spray up to rosette stage. |
| Wild Turnip | 730mL | 660mL# | 460mL | 1.35L | 460-730mL | 600mL | Spray up to rosette stage. |

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application.

DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING

- Use 30-120 litres of water per hectare.
- Use minimum of 170 litres water per hectare for LINSEED.

AERIAL SPRAYING

- Use 10 - 90 litres water per hectare.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank and Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with Metsulfuron 600, Banvel*, Apparent Bromoxynil 200, Flowable Diuron, Apparent Clopyralid 300, Sertin*, Apparent Paraquat 250, Flowable Terbutryn, Tordon* and Tribunil*. The effectiveness of Imidan*, Le Mat*, Metasystox* and dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraybuff* or Spraymate* L1-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.