

Product Name: CYDECTIN ORAL DRENCH FOR SHEEP
 APVMA Approval No: 45738/122061



Label Name:	CYDECTIN ORAL DRENCH FOR SHEEP
-------------	--------------------------------

Signal Headings:	<p>CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
------------------	--

Constituent Statements:	1 g/L MOXIDECTIN
-------------------------	------------------

Claims:	<p>For the treatment and control of moxidectin sensitive gastrointestinal parasites (including benzimidazole and/or levamisole resistant strains), lungworm and itchmite of sheep.</p> <p>CYDECTIN ORAL DRENCH FOR SHEEP contains moxidectin, a second generation macrocyclic lactone endectocide. It is effective against internal and external parasites sensitive to this family including the following parasites of sheep:</p> <p>Mature (adult) and immature (L4): Haemonchus contortus - Barbers pole worm Trichostrongylus spp. - Black scour worm Trichostrongylus axei - Stomach hair worm Trichostrongylus colubriformis Teladorsagia (Ostertagia) spp. - Small brown stomach worm Teladorsagia (Ostertagia) circumcincta Cooperia spp. - Small intestinal worm Cooperia oncophora Nematodirus spp. - Thin necked intestinal worm Nematodirus battus Nematodirus filicollis Nematodirus spathiger Strongyloides papillosus - Intestinal threadworm Oesophagostomum columbianum - Nodule worm Chabertia ovina - Large mouthed bowel worm Dictyocaulus filaria - Large lungworm</p> <p>Adult nematodes: Trichostrongylus rugatus Trichostrongylus vitrinus Cooperia curticei Nematodirus abnormalis Nematodirus helvetianus</p>
---------	---

Teladorsagia (Ostertagia) trifurcata
Oesophagostomum venulosum - Large bowel worm
Trichuris ovis

Ectoparasites:
Psorobia ovis - Itchmite

PERSISTENT ACTIVITY:

CYDECTIN ORAL DRENCH FOR SHEEP when used as recommended prevents:

- reinfection with Haemonchus contortus worms for at least 14 days and pasture contamination by H. contortus eggs for at least 30 days after a single treatment;
- reinfection with Teladorsagia (Ostertagia) spp. worms for at least 14 days and pasture contamination by Teladorsagia (Ostertagia) spp. eggs for at least 35 days after a single treatment.

CYDECTIN ORAL DRENCH FOR SHEEP is effective against benzimidazole and levamisole resistant strains of H. contortus, T. colubriformis and Tela. circumcincta.

CYDECTIN has shown efficacy against a number of ivermectin resistant strains of Haemonchus contortus and Teladorsagia (Ostertagia) spp. However, as CYDECTIN shares a similar mechanism of action to the avermectins, it SHOULD NOT be used where avermectin resistance is present, except where specifically recommended by an advisor.

RESISTANCE AND WORM CONTROL PROGRAMS

Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

Net Contents:

100mL
5L
10L
10.5L
15L
15.75L
20L
21L

Directions for Use:

Restrains:

DO NOT USE in ewes which are producing or may in the future produce milk or milk products for human consumption.

Contraindications:

Precautions:

CYDECTIN ORAL DRENCH FOR SHEEP is not recommended for use in goats.

Side Effects:

Dosage and Administration:

CYDECTIN ORAL DRENCH FOR SHEEP is ready to use. Administer 1 mL per 5 kg body weight using any standard drenching equipment. Check dose rates and equipment before drenching commences.

DOSE RATE

1 mL/5 kg body weight to give a dose of 0.2 mg moxidectin/kg body weight.

	<p>[DOSAGE TABLE attached]</p> <p>Animals heavier than 75 kg to be dosed at 1 mL per 5 kg. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.</p> <p>This section contains file attachment.</p>
--	---

General Directions:	<p>CYDECTIN ORAL DRENCH FOR SHEEP has a wide margin of safety when used as recommended and is readily accepted by sheep.</p> <p>CYDECTIN ORAL DRENCH FOR SHEEP is safe to use concurrently with other treatments including mineral supplements, fluke treatments and vaccines.</p>
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 7 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI):</p> <p>DO NOT USE less than 7 days before slaughter for export. Before using this product confirm the current ESI from either Virbac (Australia) Pty Ltd on 1800 242 100 or the APVMA website (www.apvma.gov.au/residues).</p>
---------------	---

Safety Directions:	<p>Poisonous if swallowed. May irritate the nose and throat and will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>Additional information is listed in the Safety Data Sheet (SDS).</p>
-------------------------	---

Environmental Statements:	<p>ENVIRONMENTAL PROTECTION:</p> <p>Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
---------------------------	--

Disposal:	<p>Containers less than or equal to 1L: Dispose of container by wrapping with paper and putting in garbage.</p> <p>Containers greater than 1L: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or</p>
-----------	--

similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage:

Store below 30°C (room temperature). Protect from light.

CYDECTIN ORAL DRENCH FOR SHEEP

Dosage and Administration – Dosage Table

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15