



DIRECTIONS FOR USE:

Contraindications:

This product should not be used when selenium intake from pasture is high or selenium is provided by other means (such as pasture top dressing, vaccine, pellets or selenium drenches) or if blood selenium levels at treatment are high. Users can determine selenium status by consulting a veterinarian.

Precautions:

Selenium can have toxic effects, particularly in sheep with liver damage that graze in areas where plants that contain pyrrolizidine alkaloids are prevalent.

Dosage and Administration:

Closicare[®] plus Selenium dose rate is 1 mL / 5 kg bodyweight. Closicare[®] plus Selenium is ready to use and does not require mixing. Closicare[®] plus Selenium can be used at the recommended doses in sheep of all ages including pregnant ewes.

Dose Volume Table:

| Bodyweight Kg | Dose Volume mL | Sheep Treated 20 Litre |
|------------------|-------------------|---------------------------|
| Up to 15 | 3 | 6667 |
| 16 - 20 | 4 | 5000 |
| 21 - 25 | 5 | 4000 |
| 26 - 30 | 6 | 3333 |
| 31 - 35 | 7 | 2857 |
| 36 - 40 | 8 | 2500 |
| 41 - 45 | 9 | 2222 |
| 46 - 50 | 10 | 2000 |
| 51 - 55 | 11 | 1818 |
| 56 - 60 | 12 | 1667 |
| 61 - 65 | 13 | 1538 |
| 66 - 70 | 14 | 1429 |
| 71 - 75 | 15 | 1333 |

Sheep in excess of 75 kg to be dosed at 1 mL per 5 kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by liveweight in the group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Indications:

Closicare[®] plus Selenium contains Closantel, a member of the Salicylanilides group of drenches. It is effective against sensitive strains of the following internal parasites:

- adult and immature Barber's pole worms (*Haemonchus contortus*); Closicare[®] plus Selenium provides sustained protection against re-infestation of Barber's pole worm in sheep for 4 weeks after dosing.
- adult and immature liver fluke (*Fasciola hepatica*).

including six week old stages.

all stages of nasal bots (*Oestrus ovis*) in sheep. Selenium supplementation will prevent white muscle disease. Treat ewes 1 month before lambing and treat lambs at marking. Where selenium responsive ill thrift or selenium responsive scouring are a problem, treat lambs at marking and again 2 months later. Consult your veterinarian to have blood samples taken to assess the selenium status of your flock.

Resistance may develop to any drench. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: NIL

SAFETY DIRECTIONS:

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear rubber gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

Additional information is listed in the Material Safety Data Sheet.

VIRBAC (AUSTRALIA) PTY. LIMITED

15 Pritchard Place, Peakhurst NSW 2210

Free call Customer Service Line: 1800 009 847

www.virbac.com.au

DISPOSAL:

Triple or pressure rinse container including exterior and cap. Dispose of rinsate and any undiluted chemical according to State legislative requirements. Container may be recycled at a drumMUSTER collection site if it is clean, dry and the drumMUSTER logo is visible. Provide cap separately.

STORAGE:

Store below 30°C (Room temperature). Store in original container. Reseal after use.

® Registered Trademark of Virbac

(B): CODE: 1P0271-4

EXP:

APVMA: 50323/20L0806



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Closicare[®]

plus Selenium

Sustained Action Oral Anthelmintic
and Flukicide for Sheep

ACTIVE CONSTITUENTS:

37.5 g/L CLOSANTEL

1.2 g/L SODIUM SELENATE

(equivalent to 0.5 g/L SELENIUM)



For the sustained control of Barber's Pole worm (*Haemonchus contortus*) and control of Liver Fluke (*Fasciola hepatica*) mature and immature - including six weeks old stages, and Nasal Bot (*Oestrus ovis*) and aids in control of selenium responsive conditions in Sheep.

20 L

Virbac
ANIMAL HEALTH

