



Product Name: CLOSICARE PLUS SELENIUM SUSTAINED ACTION ORAL ANTHELMINTIC AND

FLUKICIDE FOR SHEEP

APVMA Approval No: 50323 / 119940

CLOSICARE PLUS SELENIUM SUSTAINED ACTION ORAL ANTHELMINTIC AND Label Name: FLUKICIDE FOR SHEEP

Signal Headings: **POISON**

KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent 37.5 g/L CLOSANTEL

Statements: 1.2 g/L SODIUM SELENATE (equivalent to 0.5 g/L SELENIUM)

Claims:

For the sustained control of Barbers Pole worm (Haemonchus contortus) and control of Liver Fluke (Fasciola hepatica) mature and immature - including six weeks old stages, and Nasal Bot (Oestrus ovis) and aids in control of selenium responsive conditions in sheep.

Indications:

Closicare plus Selenium contains Closantel, a member of the Salicylanilides group of drenches. It is effective against sensitive strains of the following internal parasites:

- adult and immature Barber's pole worms (Haemonchus contortus).

Closicare plus Selenium provides sustained protection against re-infestation of Barber's pole worm in sheep for 4 weeks after dosing.

- adult and immature liver fluke (Fasciola hepatica) including six week old stages.
- all stages of nasal bots (Oestrus ovis) in sheep.

Selenium supplementation will prevent white muscle disease. Treat ewes 1 months before lambing and treat lambs at marking. Where selenium responsive ill thrift or selenium responsive scouring are a problem, treat lambs at marking and again 2 months later. Consult your veterinarian to have blood samples taken to assess the selenium status of your flock.

Resistance may develop to any drench. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

Net Contents: 1L 5L

20L

	25L
Directions for Use:	
Restraints:	DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
Contraindications:	This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.
Precautions:	Selenium can have toxic effects, particularly in sheep with liver damage that graze in areas where plants that contain pyrrolizidine alkaloids are prevalent.
Side Effects:	
Danamand	Olasiaana who Calarium daaa wata ia 4 wal / 5ba hada wainkt Olasiaana who Calarium
Dosage and Administration:	Closicare plus Selenium dose rate is 1 mL / 5kg body weight. Closicare plus Selenium is ready to use and does not require mixing. Closicare plus Selenium can be used at the recommended doses in sheep of all ages including pregnant ewes.
	[Dose Volume Table_Closicare plus Se]
	Sheep in excess of 75 kg to be dosed at 1 mL per 5 kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by liveweight in the group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.
	This section contains file attachment. File Name: Dose Volume Table_Closicare plus Se.docx File Size: 13095 bytes
General Directions:	
Withholding Periods:	WITHHOLDING PERIODS: MEAT: DO NOT USE less than 28 days before slaughter for human consumption. MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 60 days before slaughter for export. Before using this product, confirm the current ESI from Virbac (Australia) Pty. Ltd. on 1800 242 100 or the APVMA website (www.apvma.gov.au/residues).
Safety Directions:	Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear rubber gloves. Wash hands after use. After each days use, wash gloves.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
First Aid Warnings:	
Additional User Safety:	Additional information is listed in the Safety Data Sheet (SDS).
Environmental Statements:	
Disposal:	Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30°C (room temperature). Store in original container. Reseal after use.

CLOSICARE PLUS SELENIUM

Dose Volume Table:

Bodyweight	Dose Volume
Kg	mL
Up to 15	3
16-20	4
21-25	5
26-30	6
31-35	7
36-40	8
41-45	9
46-50	10
51-55	11
56-60	12
61-65	13
66-70	14
71-75	15