

Alben 20L  
 DIMENSIONS: 60 x 150, 430 x 270  
 DATE: 13/09/06  
 INKS: 4 COLOUR PROCESS

Label image provided by

  
**Infopest**  
 infopest@dpi.qld.gov.au



**Alben**

Broad Spectrum Anthelmintic  
 for Sheep, Lambs and Goats

**■ DIRECTIONS FOR USE:**

**SHAKE WELL BEFORE USE.** A representative sample of animals should be weighed before treatment.

**Dosage and Administration:**

**Sheep, Lambs & Goats:**

3.8 mg/kg Albendazole or 1 mL per 5 kg bodyweight for roundworms, lungworms and tapeworms. 4.75 mg/kg Albendazole or 1 mL per 4 kg bodyweight for adult liver fluke.

Bodyweight Kg	Roundworms, Lungworms & Tapeworms mL	20 L Container treats	Adult Liver Fluke Roundworms, Lungworms & Tapeworms mL	20 L Container treats
Under 15	3	6600	4.0	5000
16-20	4	5000	5.0	4000
21-25	5	4000	6.5	3076
26-30	6	3333	7.5	2666
31-35	7	2857	9.0	2222
36-40	8	2500	10.0	2000
41-45	9	2222	11.5	1739
46-50	10	2000	12.5	1600
51-55	11	1818	14.0	1428
56-60	12	1666	15.0	1333
61-65	13	1538	16.5	1212
66-70	14	1428	17.5	1142
71-75	15	1333	19.0	1052

Sheep and goats in excess of 75 kg to be dosed at 1 mL per 5 kg bodyweight for roundworms, lungworms and tapeworms or 1 mL per 4 kg bodyweight for adult liver fluke. Where adult liver fluke infection exists concurrently with roundworm, tapeworm and lungworm infections the higher dose rate should be used. Animals should be weighed. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs; bucks, does, kids). Do not underdose. Check accuracy of drenching gun before and during use. Rinse drenching guns after use.

**■ MEDICINAL CLAIM:**

**SHEEP, LAMBS & GOATS:**

For the control of Benzimidazole-Sensitive mature and immature Gastrointestinal Roundworms, Large Lungworms, Tapeworms and aids in control of Adult Liver Fluke in sheep, lambs and goats.

- (*Haemonchus contortus*) Barber's pole worm;
- (*Trichostrongylus axei*) Stomach hair worm;
- (*Ostertagia/Teladorsagia* spp) Small brown stomach worm;
- (*Trichostrongylus* spp) Black scour worm;
- (*Cooperia* spp) Small intestinal worm;
- (*Nematodirus* spp) Thin necked intestinal worm;
- (*Oesophagostomum columbianum*) Nodule worm;
- (*Oesophagostomum venulosum*) Large bowel worm;

(*Chabertia ovina*) Large mouthed bowel worm;  
 (*Dictyocaulus filaria*) Large lungworm;  
 (*Moniezia expansa*) Tapeworm;  
 (*Fasciola hepatica*) Liver Fluke.  
 Ovicidal: reduces worm egg and fluke egg hatching. For this to be of benefit, sheep should not be placed on clean pasture until 6 hours after drenching. Resistance may develop to any drench. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

**■ WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 10 days before slaughter for human consumption.**

**MILK: DO NOT USE in animals producing milk for human consumption.**

**■ TRADE ADVICE:** Export Slaughter Interval (ESI): DO NOT USE less than 10 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from either the reseller or the APVMA website [www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)

**■ SAFETY DIRECTIONS:**

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

**■ FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

Additional Information is in the Material Safety Data Sheet.

**VIRBAC (AUSTRALIA) PTY LIMITED**

15 Pritchard Place, Peakhurst NSW 2210

Freecall Customer Service Line: 1800 009 847

[www.virbac.com.au](http://www.virbac.com.au)

**■ DISPOSAL:**

Triple or pressure rinse container including exterior and cap. Dispose of rinsate and any undiluted chemical according to State legislative requirements. Container may be recycled at a drumMUSTER collection site if it is clean, dry and the drumMUSTER logo is visible. Provide cap separately.

**■ STORAGE:**

Store below 30°C (Room Temperature) with top secure - Avoid Freezing.

CODE: 1P0099-3  
 APVMA: 40645/20L/0806

(B):

EXP:



9 512214 207009

**CAUTION**  
 KEEP OUT OF REACH OF CHILDREN  
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
 FOR ANIMAL TREATMENT ONLY

**Alben**

Broad Spectrum Anthelmintic  
 for Sheep, Lambs and Goats

ACTIVE CONSTITUENT:  
 19 g/L ALBENDAZOLE




For the control of Benzimidazole-sensitive mature and immature Gastrointestinal Roundworms, Large Lungworms, Tapeworms and aids in control of Adult Liver Fluke in sheep, lambs and goats.

**Alben**

Broad Spectrum Anthelmintic  
 for Sheep, Lambs and Goats

 **Ovicidal**

 **Registered  
 for goats**

 **Aids in control  
 of adult liver  
 fluke**

 **Controls  
 tapeworm**



**Virbac**  
 ANIMAL HEALTH

20 L