

APVMA Approval No: 47057/120978

Side Effects:



Label Name: SELPOR SELENIUM POUR-ON FOR CATTLE Signal Headings: DANGEROUS POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING Constituent 5 g/L SELENIUM (as sodium selenate) Statements: Also contains: 300 g/L Diethylene glycol monobutyl ether (solvent) Claims: A selenium supplement for cattle in selenium deficient areas. Net Contents: 1L 2.5L 5L Directions for Use: Restraints: Contraindications: This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) or if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian. Precautions: Do not use if rain is imminent. Do not use in cattle with light coats intended for showing, as a brown discoloration may occur. The treatment of wet animals may reduce efficacy. When handling animals in the days immediately after treatment, wear PVC gloves.

Dosage and Administration:	SHAKE WELL BEFORE USE. To be used by direct application on the backline of cattle. Pour the required quantity along the backline from between the shoulders to the tail, using an applicator. Repeat doses should be made on the basis of diagnosed need only. The dose is 2 mL per 50 kg or 20 mL per adult cattle. This section contains file attachment.	
General Directions:		
Withholding Periods:	MEAT: Zero (0) days MILK: Zero (0) days	
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days. Before using this product, confirm the current ESI from Virbac (Australia) Pty. Limited on 1800 242 100 or the APVMA website (www.apvma.gov.au/residues).	
Safety Directions:	Poisonous if absorbed by skin contact or swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.	
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.	
First Aid Warnings:		
Additional User Safety:	Additional information is listed in the Safety Data Sheet (SDS).	
Environmental Statements:		
Disposal:	Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection	

or similar container management program site. The cap should not be replaced, but may be recycled separately with the container.

Storage:	Store below 30 °C (room temperature).
----------	---------------------------------------

Selpor Selenium Pour-on for Cattle

Dosage Table:

Liveweight kg	Dose Volume (mL)
Up to 50	2
51 - 100	4
101- 150	6
151 - 200	8
201 - 250	10
251 - 300	12
301 - 350	14
351 - 400	16
401 - 450	18
451 and over	20