



**Company Name:** MAVLAB PTY. LTD.  
**Product Name:** FIDO'S CLOSASOLE 10KG BROADSPECTRUM WORMER FOR DOGS & CATS

**APVMA Approval No:** 37422/103541



<b>Label Name:</b>	FIDO'S CLOSASOLE 10KG BROADSPECTRUM WORMER FOR DOGS & CATS
<b>Signal Headings:</b>	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
<b>Constituent Statements:</b>	Each tablet contains: 1000 mg NICLOSAMIDE and 41.8 mg LEVAMISOLE
<b>Claims:</b>	For the control of roundworm (Toxocara & Toxascaris), Hookworm (Uncinaria & Ancylostoma) and tapeworm (Dipylidium & Taenia) but not hydatid (Echinococcus granulosus) in dogs and cats. Does not control heartworm or hydatids in dogs. If worm problem persists, contact a veterinarian.
<b>Net Contents:</b>	10 Tablets 100 Tabs
<b>Directions for Use:</b>	Cats: Tapeworms: Treat every 3 months Roundworms & hookworm: Treat at 6, 8 and 12 weeks - thereafter every 3 months Pregnant cats: Treat at mating, before birth of kittens, then every 3 months Dogs: Tapeworms: Treat every 3 months (To prevent re-infestation with Dipylidium spp a flea control regime is essential) Roundworm & hookworm: Treat at 2, 4, 8 and 12 weeks - thereafter every 3 months Whipworm: Treat every 6 to 8 weeks after 3 months of age Pregnant bitches: Treat at mating, before whelping and then every 3 months
<b>Side Effects:</b>	Side effects in a small proportion of animals are vomiting, depressed appetite and unsteady gait. If unsteady gait or depressed appetite occurs, discontinue administration of the drug. To prevent vomiting, administer with food.

<b>Dosage and Administration:</b>	Adult Dogs & Cats: 1 tablet/10 kg Bodyweight (BW), once every 3 months throughout life. Bitches And Queens: 1 tablet/10 Kg BW, 2 treatments (at an interval of 14 days) during the gestation and lactation periods.
<b>General Directions:</b>	Administer with a light meal or crushed in food. If used in performance animals, the regulations of the relevant authorities regarding medication should be observed. Consult your veterinarian for further advice.
<b>First Aid Instructions:</b>	If poisoning occurs, contact a doctor or Poisons Information Centre (Phone – Australia 13 11 26).
<b>First Aid Warnings:</b>	Not applicable
<b>Disposal:</b>	Dispose of empty container by wrapping with paper and putting in garbage.
<b>Storage:</b>	Store below 30°C (Room Temperature). Protect from light. Replace lid tightly after use.