

Product Name: PARAMECTIN POUR-ON FOR CATTLE
 APVMA Approval No: 50341/118710



Label Name:	PARAMECTIN POUR-ON FOR CATTLE
Signal Headings:	<p>POISON</p> <p>KEEP OUT OF REACH OF CHILDREN</p> <p>FOR ANIMAL TREATMENT ONLY</p> <p>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
Constituent Statements:	<p>5 g/L ABAMECTIN</p> <p>Also contains: 950 g/L DIETHYLENE GLYCOL MONOBUTYL ETHER (solvent)</p>

Claims:	<p>For the treatment and control of abamectin sensitive internal and external parasites of cattle.</p> <p>INDICATIONS</p> <p>PARAMECTIN Pour-On for Cattle contains abamectin, a member of the avermectin family of compounds. It provides a broad spectrum of efficacy against sensitive strains of the following adult and immature internal and external parasites of cattle. Resistance may develop to any drench.</p> <p>Adult and immature gastrointestinal roundworms Haemonchus placei - Barber's pole worm Ostertagia ostertagi - Small brown stomach worm (including inhibited L4 stage). Controls infection acquired up to 14 days after treatment. Trichostrongylus axei - Stomach hair worm Trichostrongylus colubriformis - Intestinal hair worm Cooperia oncophora, C. punctata - Small intestinal worm Nematodirus spathiger - Thin-necked intestinal worm Oesophagostomum radiatum - Nodule worm Oesophagostomum venulosum - Large bowel worm Trichuris spp. - Whip worm Strongyloides papillosus - Intestinal threadworm.</p> <p>Lungworms Dictyocaulus viviparus - Large lungworm. Controls infection acquired up to 28 days after treatment.</p> <p>Sucking Lice and Biting Lice Linognathus vituli - Long-nosed sucking louse Haematopinus eurysternus - Short-nosed sucking louse Bovicola bovis - Biting louse Solenopotes capillatus - Little blue sucking louse.</p> <p>Cattle Tick: Boophilus microplus</p>
---------	---

	<p>Buffalo Fly: <i>Haematobia irritans exigua</i>. Controls buffalo fly and protects cattle from reinfestation for up to 14 days after treatment.</p> <p>PARAMECTIN Pour-On for Cattle is not recommended in strategic dipping programmes.</p>
--	--

Net Contents:	[1 L] [2.5 L] [5 L] [10 L] [20 L]
---------------	-----------------------------------

Directions for Use:	
---------------------	--

Restrains:	The minimum retreatment interval permitted is 35 days.
------------	--

Contraindications:	This product is contraindicated for use in calves under 16 weeks of age or less than 50 kg of bodyweight.
--------------------	---

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>Dose the mob according to the heaviest animal by liveweight in the group (bulls, cows, steers, calves etc.) Do not underdose. A representative sample of animals should be weighed before treatment either with scales or a weighband. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.</p> <p>Dose: 1 mL per 10 kg bodyweight (500 µg/kg bodyweight).</p> <p>Dose table: [Attached]</p> <p>Cattle in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.</p>
----------------------------	--

General Directions:	
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 35 days before slaughter for human consumption.</p> <p>MILK: Zero (0) days.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues).</p>
---------------	---

Safety Directions:	Poisonous if absorbed by skin contact or inhaled or swallowed. Will damage eyes. May irritate the nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. When using the product wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.
First Aid Warnings:	
Additional User Safety:	
Environmental Statements:	<p>Abamectin is extremely toxic to aquatic species. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p> <p>Effect on Dung Beetles When used according to the label directions, Paramectin Pour-On for Cattle is not likely to have any unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations of the following species: <i>Onthophagus australis</i>, <i>O. granulatus</i>, <i>O. dandalu</i>, <i>O. chepara</i>, <i>Onitis alexis</i>, <i>Euoniticellus intermedius</i>, <i>E. africanus</i> and <i>Aphodius</i> spp. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.</p>
Disposal:	<p>DISPOSAL: [1 L pack size] Dispose of empty container by wrapping with paper and putting in garbage.</p> <p>[2.5 L, 5 L, 10 L, 20 L pack sizes] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
Storage:	Store below 30°C (room temperature). Store in original container. Protect from light.

Liveweight	Dose	1 L treats	2.5 L treats	5 L treats	10 L treats	20 L treats
50-75 kg	7.5 mL	133	333	666	1333	2666
76-100 kg	10 mL	100	250	500	1000	2000
101-150 kg	15 mL	66	166	333	666	1333
151-200 kg	20 mL	50	125	250	500	1000
201-250 kg	25 mL	40	100	200	400	800
251-300 kg	30 mL	33	83	166	333	666
301-350 kg	35 mL	28	71	140	285	571
351-400 kg	40 mL	25	62	125	250	500
401-450 kg	45 mL	22	55	110	222	444
451-500 kg	50 mL	20	50	100	200	400
501-550 kg	55 mL	18	45	90	181	363
551-600 kg	60 mL	16	41	80	166	333
601-650 kg	65 mL	15	38	75	153	307

Cattle in excess of 650 kg to be dosed at 5 mL per 50 kg bodyweight.