



Product Name: CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS
APVMA Approval No: 52085/118661



Label Name:	CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS
-------------	---

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	18.8 mg/mL PRAZIQUANTEL 1 mg/mL MOXIDECTIN
-------------------------	---

Claims:	<p>For the treatment and control of moxidectin sensitive gastrointestinal parasites (including benzimidazole and/or levamisole resistant strains), lungworm, itchmite and tapeworm of sheep and lambs.</p> <p>CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS is a second generation macrocyclic lactone endectocide which is effective against parasites sensitive to this family and praziquantel which is effective against tapeworm. The following parasites can be treated with this product;</p> <p>Mature (adult) and immature (L4): Moniezia expansa - Tapeworm Haemonchus contortus - Barber's pole worm Trichostrongylus spp. - Black scour worm Trichostrongylus axei - Stomach hair worm Trichostrongylus colubriformis Teladorsagia (Ostertagia) spp. Teladorsagia (Ostertagia) circumcincta Cooperia spp. - Small intestinal worm Cooperia oncophora Nematodirus spp. - Thin necked intestinal worm Nematodirus battus Nematodirus filicollis Nematodirus spathiger Strongyloides papillosus - Intestinal threadworm Oesophagostomum columbianum - Nodule worm Chabertia ovina - Large mouthed bowel worm Dictyocaulus filaria - Large lungworm</p> <p>Adult nematodes:</p>
---------	---

	<p>Trichostrongylus rugatus Trichostrongylus vitrinus Cooperia curticei Nematodirus abnormalis Nematodirus helvetianus Ostertagia trifurcata Oesophagostomum venulosum - Large bowel worm Trichuris ovis</p> <p>Ectoparasites: Psorobia ovis - Itchmite</p> <p>PERSISTENT ACTIVITY CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS when used as recommended prevents reinfection with Teladorsagia (Ostertagia) circumcincta worms for at least 14 days and pasture contamination by Teladorsagia (Ostertagia) circumcincta eggs for at least 35 days after a single treatment.</p> <p>As a result of adding tapeworm activity, this product, unlike other CYDECTIN formulations does not offer persistent activity against barber's pole worm (Haemonchus contortus). However, it is fully effective against all stages of barber's pole worm infection at the time of treatment.</p> <p>CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS is effective against benzimidazole and levamisole resistant strains of H. contortus, T. colubriformis and Tela. circumcincta. CYDECTIN has shown efficacy against a number of avermectin resistant strains of H. contortus and Ostertagia spp. However, as CYDECTIN shares a similar mechanism of action to the avermectins, it SHOULD NOT be used where avermectin resistance is present, except where specifically recommended by an advisor.</p> <p>RESISTANCE AND WORM CONTROL PROGRAMS Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce the development of resistance. It is advisable that a drench resistance test be conducted before any drench is used. Resistance may develop to any drench.</p>
--	--

Net Contents:	5L 10L 15L 15.75L 20L
---------------	-----------------------------------

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE in ewes, which are producing or may in the future produce milk or milk products for human consumption.
------------	---

Contraindications:	
--------------------	--

Precautions:	CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS is not recommended for use in goats as safety and efficacy has not been evaluated.
--------------	--

Side Effects:	
Dosage and Administration:	<p>Shake well before use to ensure re-suspension. Administer 1 mL per 5 kg liveweight using any standard drenching equipment. Check dose rates and equipment before drenching commences.</p> <p>DOSE RATE 1 mL/5 kg liveweight to give a dose of 0.2 mg moxidectin/kg and 3.76 mg praziquantel/kg liveweight.</p> <p>[Dosage table - see Attachment 1 - Dosage Table]</p> <p>Animals above 75 kg to be dosed at 1 mL per 5 kg liveweight. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.</p> <p>This section contains file attachment.</p>
General Directions:	<p>Use this product for tapeworm and roundworm control. Use CYDECTIN Oral drench for Sheep in routine worm control programs where tapeworm is not present. CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS has a wide margin of safety when used as recommended and is readily accepted by sheep. CYDECTIN PLUS TAPE ORAL DRENCH AND TAPEWORM TREATMENT FOR SHEEP AND LAMBS is safe to use concurrently with other treatments including fluke treatments and vaccines.</p>
Withholding Periods:	<p>MEAT: DO NOT ADMINISTER less than 7 days before slaughter for human consumption. MILK: DO NOT USE in ewes, which are producing or may in the future produce milk or milk products for human consumption.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. Before using this product confirm the current ESI from either Virbac (Australia) Pty. Ltd. on 1800 242 100 or the APVMA website (www.apvma.gov.au/residues).</p>
Safety Directions:	<p>Poisonous if swallowed. May irritate the nose and throat and will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.</p>
First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
First Aid Warnings:	

Additional User Safety:	Additional information is listed in the Safety Data Sheet (SDS).
-------------------------	--

Environmental Statements:	ENVIRONMENTAL PROTECTION Extremely toxic to aquatic species. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.
---------------------------	---

Disposal:	<p>Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container.</p> <p>If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------	--

Storage:	Store below 25 °C (air conditioning) and protect from sunlight.
----------	---

DOSAGE GUIDE

5L Pack

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15

10L Pack

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15

15 L Pack

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15

15.75 L Pack

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15

20 L Pack

Body Weight (kg)	Dose Volume (mL)
10	2
11 - 15	3
16 - 20	4
21 - 25	5
26 - 30	6
31 - 35	7
36 - 40	8
41 - 45	9
46 - 50	10
51 - 55	11
56 - 60	12
61 - 65	13
66 - 70	14
71 - 75	15