

Product Name: Y-TEX AGRESSOR CATTLE EAR TAGS
APVMA Approval No: 62199/119798



Label Name:	Y-TEX AGRESSOR INSECTICIDAL CATTLE EAR TAGS
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	80 g/kg ABAMECTIN 180 g/kg PIPERONYL BUTOXIDE
Claims:	Two tags/head treatment for up to 16 weeks control of abamectin-susceptible Buffalo Flies* on beef and dairy cattle (including lactating dairy cows). *Including organophosphate-resistant and synthetic pyrethroid-resistant strains. Resistance may develop to any chemical.
Net Contents:	20 tags - each 9 g [180 g NET weight] 100 tags - each 9 g [900 g NET weight] 500 tags - each 9 g [4.5 kg NET weight] 1000 tags - each 9 g [9.0 kg NET weight] 1300 tags - each 9 g [11.7 kg NET weight] It is illegal to sell individual tags separately
Directions for Use:	
Restrains:	Remove tags prior to slaughter.
Contraindications:	This product is contraindicated for use in calves less than 16 weeks of age or less than 50 kg bodyweight.
Precautions:	

Side Effects:	
Dosage and Administration:	<p>Animal and Pest: Beef and dairy cattle (including lactating dairy cows), Buffalo Flies (<i>Haematobia irritans exigua</i>).</p> <p>State: All states and territories.</p> <p>Rate: Two (2) tags per animal (one in each ear).</p> <p>Apply only when fly numbers exceed acceptable levels. Removal of tags after 16 weeks will ensure that Buffalo Fly populations are not exposed to sub-lethal doses which may increase the chance of resistance.</p> <p>Replace as necessary. Remove tags with tag remover. The level and duration of control may be reduced under high fly pressure. The whole herd is to be treated concurrently.</p> <p><Dosage and Administration Table></p> <p>This section contains file attachment.</p>
General Directions:	<p>Y-TEX Agressor Insecticidal Cattle Ear Tags will give effective control of Buffalo Flies if used regularly on all animals in the group (over 16 weeks of age).</p> <p>The treatment regime with a periodic rotation program will reduce the potential for insecticide resistance to develop in Buffalo Flies.</p> <p>Application: Apply tags with the Y-TEX UltraTagger® (Figures 1 to 4). Figure 1 <drawing> Seat male button firmly on pin, press down and give a slight twist.</p> <p>Figure 2 <drawing> Place female tag firmly under clip by depressing lever. Collar on tag must be pointing away from pin.</p> <p>Figure 3 <drawing> Position male stud between the second and third radial cartilage on the front of the ear with the insecticidal ear tag (female tag) on the back of the ear. Do not allow shaft of the male button to penetrate any cartilage rib or blood vessels or ear damage may result.</p> <p>Figure 4 <drawing> Squeeze handles of applicator together. When properly positioned, the tag should hang as shown.</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p>
Withholding Periods:	<p>MEAT: Zero (0) days</p> <p>MILK: Zero (0) days</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT slaughter for export less than 42 days after application. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from Landmark Operations Limited 1800 448 892 or the APVMA website www.apvma.gov.au/residues.</p>

Safety Directions:	Harmful if sucked. When using the product, wear rubber gloves. Wash hands after use.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
First Aid Warnings:	
Additional User Safety:	<p>ADDITIONAL USER SAFETY INFORMATION:</p> <p>Additional information is listed in the Safety Data Sheet.</p>
Environmental Statements:	ENVIRONMENTAL PROTECTION: Very toxic to aquatic life. DO NOT contaminate streams, rivers or waterways with the tags or used containers.
Disposal:	<p>[20, 100 tags]: Dispose of container by wrapping with paper and putting in garbage.</p> <p>[500, 1000, 1300 tags]: Break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or spent tags.</p>
Storage:	Store below 30°C (Room temperature). Store in original container in a dry, cool place. DO NOT store in direct sunlight.

Dosage and Administration

Animal and Pest	State	Rate	Critical Comments
Beef and Dairy Cattle (including lactating dairy cows) Buffalo Flies (<i>Haematobia irritans exigua</i>)	All States and Territories	Two tags per animal (One tag per ear)	Apply only when fly numbers exceed acceptable levels. Replace as necessary. Remove tags with tag remover. Removal of tags after 16 weeks will ensure that Buffalo Fly populations are not exposed to sublethal doses which may increase the chance of resistance. The level and duration of control may be reduced under high fly pressure. The whole herd is to be treated concurrently.