

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



ROUNDWORM, LIVER FLUKE AND EXTERNAL PARASITICIDE FOR CATTLE

5 mg/mL ABAMECTIN, 300 mg/mL TRICLABENDAZOLE

For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

15 LITRES

©AVOMECH and the cattle head logo are registered trademarks of the Boehringer Ingelheim Group. LF18281



**Boehringer
Ingelheim**

AVOMECS PLUS Pour-On contains Abamectin, a member of the macrocyclic lactones family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains Triclabendazole, a member of the benzimidazoles family of chemicals, effective against liver fluke. Resistance may develop to any chemical.

INDICATIONS

For the treatment and control of adult and immature stages of:

Gastro-intestinal roundworms

Barber's Pole Worm (*Haemonchus placei*), Small Brown Stomach Worm (*Ostertagia ostertagi*) (including inhibited stages), Small Intestinal Worm (*Cooperia oncophora*) (*Cooperia punctata*), Stomach Hair Worm (*Trichostrongylus axei*), Hookworm (*Bunostomum phlebotomum*), Large Bowel Worm (*Oesophagostomum radiatum*), Whipworm (*Trichuris* spp.), Thin Necked Intestinal Worm (*Nematodirus spathiger*), Intestinal Threadworm (*Strongylides papillosus*)

Lungworms (*Dictyocaulus viviparus*)

Liver Fluke (*Fasciola hepatica*) including adult (12 week old), immature (6 week old) and early immature (4 week old) larvae

Sucking and Biting Lice

Long Nosed Sucking Lice (*Linognathus vituli*), Short Nosed Sucking Lice (*Haematopinus eurysternus*), Biting Lice (*Bovicola bovis*), Little Sucking Lice (*Solenopotes capillatus*).

Also aids in the control of:

Buffalo Fly (*Haematobia irritans exigua*)

Cattle Tick (*Boophilus microplus*)

(Not recommended in strategic dipping programs for cattle tick control).

DIRECTIONS FOR USE

Restraints

DO NOT USE in dairy animals producing or which will in the future be producing milk for human consumption.

This product is contraindicated for use in calves under 50 kg bodyweight.

Dosage and Administration

AVOMECS PLUS Pour-On dose rate is 1 mL/10 kg body weight.

Apply along the backbone of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

Live weight (kg)	Dose Volume (mL)	15 Litre Pack treats
51-75	7.5	2000
76-100	10	1500
101-125	12.5	1200
126-150	15	1000
151-175	17.5	857
176-200	20	750
201-250	25	600
251-300	30	500
301-350	35	428
351-400	40	375
401-450	45	333
451-500	50	300
501-550	55	272
551-600	60	250
601-650	65	230

Over 650 kg dose at rate of 1 mL/10 kg body weight. A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

AVOMECS PLUS Pour-On is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product. A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions.

Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian. After use clean all equipment with cooking oil. Do not use water.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

MEAT: DO NOT USE less than 49 days before slaughter for human consumption.

Beef calves born to cows that were treated with AVOMECS PLUS POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the dam.

MILK: DO NOT USE in dairy animals producing or which will in the future be producing milk for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT use less than 140 days before slaughter for export. This ESI was correct at the time of label approval. Before using this product confirm the current ESI from the registrant (1800 808 691) or the APVMA website (www.apvma.gov.au/residues)

Beef calves born to cows that were treated with AVOMECS PLUS POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the dam.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product in eyes, first quickly, gently and completely **blot or wipe** chemical from eyes then flush with water. Wash hands after use. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing, quickly and gently **blot or wipe** away chemical with a tissue or cloth, preferably impregnated with cooking oil, then wash skin thoroughly using soap and water. Do not use water first.

ADDITIONAL USER SAFETY INFORMATION

Additional information is listed in the Safety Data Sheet.

ENVIRONMENTAL PROTECTION

Abamectin is extremely toxic to aquatic species.

Do not contaminate dams, rivers, streams or waterways with the chemical or used containers.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo. Triple rinse containers before disposal. Dispose of rinsate and any undiluted chemical according to State legislative requirements. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. Do not burn empty containers or product.

STORAGE

Store below 30°C (Room Temperature).

Additional GHS statements required by Safe Work Australia: Harmful if inhaled. Causes damage to organs through prolonged or repeated exposure. Do not breathe fumes, mists, vapours or spray. Do not get on clothing. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. If swallowed, rinse mouth. Do NOT induce vomiting. If inhaled, remove victim to fresh air and keep at rest in a position comfortable for breathing. If skin irritation occurs or eye irritation persists, get medical advice. Protect from sunlight. Store in a closed container, in a dry place.

Boehringer Ingelheim Animal Health Australia Pty Ltd
Level 1, 78 Waterloo Road,
North Ryde NSW 2113



For more information, contact **Boehringer Ingelheim** customer solutions on **1800 808 691**.

APVMA No: 62708/15/L0609

(B)

EXP

LB18280



Product Name:

AVOMECS PLUS POUR-ON 15L - Back Label