

100 % of final size = 200 mm x 175 mm

200 mm

## CAUTION


KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

# Strategik Mini-Dose

WORMING DRENCH FOR CATTLE

ACTIVE CONSTITUENT: 112.5 g/L ALBENDAZOLE



For the treatment and control in cattle of  
susceptible mature and immature  
gastrointestinal roundworms, (including  
inhibited Type II *Ostertagia* larvae), large  
lungworms and tapeworms; to aid in  
the control of adult liver fluke; and  
to reduce the output of viable worm  
and fluke eggs.

20L



**jurox**

407761

100 % of final size = 200 mm x 175 mm

200 mm

# Strategik Mini-Dose Worming Drench for Cattle

#### EFFICACY SPECTRUM:

STRATEGIK MINI-DOSE Worming Drench for Cattle is intended for use against benzimidazole susceptible worms of the following species:

#### Roundworms (mature and immature)

- *Haemonchus placei* - Barber's pole worm
- *Ostertagia ostertagi* - Small brown stomach worm
- *Trichostrongylus axei* - Stomach hair worm
- *Trichostrongylus* spp - Black scour worm
- *Cooperia* spp - Small intestinal worm
- *Bunostomum* spp - Hookworm
- *Oesophagostomum radiatum* - Nodule worm
- *Nematodirus* spp - Thin-necked intestinal worm
- *Strongyloides* spp - Intestinal threadworm
- *Ostertagia* spp - Inhibited Type II larvae

#### Lungworms

- *Dictyocaulus viviparus* - Large lungworm

#### Tapeworms

- *Moniezia* spp - Heads and segments

#### Liver Fluke (adults)

- *Fasciola hepatica*

#### Ovicular

Reduces the output of viable worm and fluke eggs.

Note: Worm parasites may develop resistance to drenches. It is advisable that an anthelmintic resistance test be carried out before any anthelmintic is used. This drench is not effective against worms resistant to benzimidazole type drenches. If resistance to this drench is suspected, seek veterinary confirmation and notify manufacturer.

STRATEGIK MINI-DOSE Worming Drench for Cattle aids in the control of adult liver fluke and reduces the output of viable fluke eggs. Contamination of pasture will be reduced by avoiding grazing recognised fluke areas for 24 hours after dosing.

A second treatment with STRATEGIK MINI-DOSE Worming Drench for Cattle one month after the first will assist to prevent contamination of pastures and will minimise chances of reinfection. To gain maximum benefit from the reduction in the output of viable worm and fluke eggs, avoid placing treated stock on clean pastures for a minimum of 4 hours after dosing.

#### DIRECTIONS FOR USE:

**DO NOT ADMINISTER to cattle producing milk for human consumption.**

#### SHAKE CONTAINER WELL BEFORE USE.

Administer orally, and check accuracy of drenching equipment before and during use. Clean drenching equipment thoroughly before and after use. Dose rate should be based on heaviest cattle in the mob. Cattle should be weighed either with scales or weighband.

#### Dose Rate: CATTLE

Roundworms, Lungworms, Tapeworms: 3 mL per 45 kg bodyweight.

Roundworms, Lungworms, Tapeworms and Adult Liver Fluke: 4 mL per 45 kg bodyweight.

#### DOSE VOLUME TABLE

Bodyweight	Roundworms, Lungworms, Tapeworms		Roundworms, Lungworms, Tapeworms, Adult Liver Fluke	
	Dose	20 L treats (head)	Dose	20 L treats (head)
0 - 45 kg	3 mL	6666	4 mL	5000
46 - 90 kg	6 mL	3333	8 mL	2500
91 - 135 kg	9 mL	2222	12 mL	1666
136 - 180 kg	12 mL	1666	16 mL	1250
181 - 225 kg	15 mL	1333	20 mL	1000
226 - 270 kg	18 mL	1111	24 mL	833
271 - 315 kg	21 mL	952	28 mL	714
316 - 360 kg	24 mL	833	32 mL	625
361 - 405 kg	27 mL	740	36 mL	555
406 - 450 kg	30 mL	666	40 mL	500
451 - 495 kg	33 mL	606	44 mL	454
496 - 540 kg	36 mL	555	48 mL	416
541 - 585 kg	39 mL	512	52 mL	384
586 - 630 kg	42 mL	476	56 mL	357
631 - 675 kg	45 mL	444	60 mL	333

Cattle in excess of 675 kg to be dosed at the appropriate rate shown before the Dose Volume Table.

#### WITHHOLDING PERIODS:

**MEAT: DO NOT ADMINISTER less than 10 days before slaughter for human consumption.**

**MILK: DO NOT ADMINISTER to cattle producing milk for human consumption.**

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT SLAUGHTER for export for 10 days after treatment.**

**SAFETY DIRECTIONS:** Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre, phone 131126.

**DISPOSAL:** Triple rinse empty container. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to point of supply. If not recycling, break, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose and clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORE below 30°C (room temperature). Store in the original container tightly closed, in a safe place away from foods.

APVMA Approval No. 48119/20L/0306

(B.)

EXP.

40776

Jurox Pty Ltd  
85 Gardiners Road  
Rutherford NSW 2320  
Infoline 1800 023 312

9 338094 000000 >