



Product Name: STRATEGIK MINI-DOSE WORMING DRENCH FOR CATTLE
APVMA Approval No: 48119/116495



Label Name:	STRATEGIK MINI-DOSE WORMING DRENCH FOR CATTLE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 112.5 g/L ALBENDAZOLE
Claims:	<p>For the treatment and control in cattle of susceptible mature and immature gastrointestinal roundworms (including inhibited Type II <i>Ostertagia</i> larvae), large lungworms and tapeworms; to aid in the control of adult liver fluke; and to reduce the output of viable worm and fluke eggs.</p> <p>EFFICACY SPECTRUM STRATEGIK MINI-DOSE Worming Drench for Cattle is intended for use against benzimidazole susceptible worms of the following species:</p> <p>Roundworms (mature and immature)</p> <ul style="list-style-type: none">• <i>Haemonchus placei</i> - Barber's pole worm• <i>Ostertagia ostertagi</i> - Small brown stomach worm• <i>Trichostrongylus axei</i> - Stomach hair worm• <i>Trichostrongylus</i> spp. - Black scour worm• <i>Cooperia</i> spp. - Small intestinal worm• <i>Bunostomum phlebotomum</i> – Hookworm• <i>Oesophagostomum radiatum</i> - Nodule worm• <i>Nematodirus</i> spp. - Thin-necked intestinal worm• <i>Strongyloides</i> spp. - Intestinal threadworm• <i>Ostertagia</i> spp. - Inhibited Type II larvae <p>Lungworms</p> <ul style="list-style-type: none">• <i>Dictyocaulus viviparus</i> - Large lungworm <p>Tapeworms</p> <ul style="list-style-type: none">• <i>Moniezia</i> spp. - Heads and Segments <p>Liver Fluke (adults)</p> <ul style="list-style-type: none">• <i>Fasciola hepatica</i> <p>Ovicidal</p> <p>Reduces the output of viable worm and fluke eggs.</p> <p>Note: Worm parasites may develop resistance to drenches. It is advisable that an anthelmintic resistance test be carried out before any anthelmintic is used. This drench is not effective against worms resistant to benzimidazole type drenches. If resistance to this drench is suspected, seek veterinary confirmation and notify manufacturer.</p>

Net Contents:	1 L 5 L 10 L 20 L
---------------	----------------------------

Directions for Use:	
---------------------	--

Restraints:	DO NOT ADMINISTER to cattle producing milk for human consumption.
-------------	---

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>SHAKE CONTAINER WELL BEFORE USE</p> <p>Administer orally, and check accuracy of drenching equipment before and during use. Clean drenching equipment thoroughly before and after use. Dose rate should be based on heaviest cattle in the mob. Cattle should be weighed either with scales or weighband.</p> <p>Dose Rates: Cattle</p> <p>Roundworms, Lungworms, Tapeworms: 3 mL per 45 kg bodyweight.</p> <p>Roundworms, Lungworms, Tapeworms, Adult Liver Fluke: 4 mL per 45 kg bodyweight.</p> <p>Dose Volume Table:</p> <p>[dose volume table attached]</p> <p>Cattle in excess of 675 kg to be dosed at the appropriate rate shown before the Dose Volume Table.</p> <p>This section contains file attachment.</p>
----------------------------	--

General Directions:	<p>STRATEGIK MINI-DOSE Worming Drench for Cattle aids in the control of adult liver fluke and reduces the output of viable fluke eggs. Contamination of pasture will be reduced by avoiding grazing recognised fluke areas for 24 hours after dosing. A second treatment with STRATEGIK MINI-DOSE Worming Drench for Cattle one month after the first will assist to prevent contamination of pastures and will minimise chances of reinfestation. To gain maximum benefit from the reduction in the output of viable worm and fluke eggs, avoid placing treated stock on clean pastures for a minimum of 4 hours after dosing.</p>
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT ADMINISTER less than 10 days before slaughter for human consumption.</p> <p>MILK: DO NOT ADMINISTER to cattle producing milk for human consumption.</p>
----------------------	---

Trade Advice:	<p>DO NOT USE less than 10 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues/).</p>
---------------	--

Safety Directions:	Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	
---------------------------	--

Disposal:	Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------	--

Storage:	Store below 30°C (room temperature). Store in the original container, tightly closed, in a safe place away from foods.
----------	--

Dose Volume Table:

	Roundworms, Lungworms, Tapeworms					Roundworms, Lungworms, Tapeworms and Adult Liver Fluke				
Body Weight	Dose	1 L Treats (head)	5 L Treats (head)	10 L Treats (head)	20 L Treats (head)	Dose	1 L treats (head)	5 L Treats (head)	10 L Treats (head)	20 L Treats (head)
0 – 45 kg	3 mL	333	1666	3333	6666	4 mL	250	1250	2500	5000
46 – 90 kg	6 mL	166	833	1666	3333	8 mL	125	625	1250	2500
91 – 135 kg	9 mL	111	555	1111	2222	12 mL	83	416	833	1666
136 – 180 kg	12 mL	83	416	833	1666	16 mL	62	312	625	1250
181 – 225 kg	15 mL	66	333	666	1333	20 mL	50	250	500	1000
226 – 270 kg	18 mL	55	277	555	1111	24 mL	41	208	416	833
271 – 315 kg	21 mL	47	238	476	952	28 mL	35	178	357	714
316 – 360 kg	24 mL	41	208	416	833	32 mL	31	156	312	625
361 – 405 kg	27 mL	37	185	370	740	36 mL	27	138	277	555
406 – 450 kg	30 mL	33	166	333	666	40 mL	25	125	250	500
451 – 495 kg	33 mL	30	151	303	606	44 mL	22	113	227	454
496 – 540 kg	36 mL	27	138	277	555	48 mL	20	104	208	416
541 – 585 kg	39 mL	25	128	256	512	52 mL	19	96	192	384
586 – 630 kg	42 mL	23	119	238	476	56 mL	17	89	178	357
631 – 675 kg	45 mL	22	111	222	444	60 mL	16	83	166	333