

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Conquest AMINE 300

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 300g/L 2, 4 -D present as the isopropylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For the control of emerged broadleaved weeds prior to sowing crops and pastures in conservation tillage situations and for selective weed control in the crops and situations as per directions for use table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals

IMPORTANT : READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 20 LITRES
(5 L, 110 L, 200 L, 1000 L)

Conquest Agrochemicals Pty Ltd.
ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696



RLP
Approved



STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry, well-ventilated area away from children, animals and food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers : Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL
1 800 033 111
24 HOURS AUSTRALIA WIDE.

Conquest Agrochemicals Pty Ltd.
ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

Batch No:
DOM:

Approval No: 55701/0214

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Conquest AMINE 300

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 300 g/L 2, 4 -D present as the isopropylamine salt

GROUP	I	HERBICIDE
-------	---	-----------

For the control of emerged broadleaved weeds prior to sowing crops and pastures in conservation tillage situations and for selective weed control in the crops and situations as per directions for use table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals

IMPORTANT: READ THIS LEAFLET BEFORE USE

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932

Suite 9, 41 Walters Drive

Herdsmen Business Park

Osborne Park WA 6017

Telephone (08) 9244 1268

Facsimile: (08) 9446 4696

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT spray if rain seems likely within 6 hours or if strong winds prevail.
DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions

1. CONSERVATION TILLAGE

SITUATION AND CROP	WEEDS	STATE	RATE/ha	CRITICAL COMMENTS
Preparatory spray for fallows and seedbeds or prior to sowing the following crops: Balansa clover, Barley, Chickpeas, Cotton, Faba beans, Field peas, Lentils, Linseed, Lucerne, Lupins, Narbon beans, Navy beans, Oats, Perennial ryegrass, Persian clover, Phalaris, Rapeseed, Rice, Safflower, Sorghum, Soybean, Subterranean clover, Sunflower, Triticale, Vetch, Wheat, White clover.	Fumitory (white), Ball Mustard, Indian Hedge Mustard, Common Sowthistle, Turnip Weed, Wild Turnip, Wild Radish	All States	660mL - 1.2L + Conquest Glyphosate 450 Herbicide (Glyphosate) and Surfactant at recommended label rates	RATE SELECTION. Use the lower rate for seedling broadleaf weeds and increase to the higher rate for broadleaf weeds more than 10cm diameter/high. Always add Conquest Glyphosate 450 Herbicide (Glyphosate) at recommended label rates. At time of application, all weeds must be actively growing and not under stress from low moisture, frost, cold, disease or waterlogging. If grazing has occurred, allow regrowth to 6-8cm before spraying and use higher rate. Always add a non-ionic surfactant (eg. Conquest Wetter 1000) in accordance with the label directions on the Conquest Glyphosate 450 Herbicide.
	Seedlings of: Australian Bindweed, Bellvine, Caltrop, New Zealand Spinach, Raspweed	NSW, ACT, Qld only		
	Ageratum (Blue top), Dock, Volunteer Lupins, Volunteer Peas, Volunteer Sunflowers, Charlock, Fumitory (Red), Medic, Paterson's Curse, Prickly Lettuce (Wild Lettuce), Saffron Thistle, Spear Thistle, Variegated Thistle	All States	900mL - 1.2L + Conquest Glyphosate 450 Herbicide (Glyphosate) and Surfactant at recommended label rates	
	Bathurst Burr, Blackberry Nightshade, Californian Burr, Horehound Seedlings, Lincoln Weed Seedlings, Marshmallow Seedlings, Sorrel Seedlings, Thornapple, Volunteer Vetch, Volunteer Safflower, Common Ice-Plant, Storksbill/Erodium Seedlings, Ivyleaf, Speedwell, Melilotus, Shepherd's Purse, Skeleton Weed (Suppression only), Ward's Weed, Wireweed Seedlings (Hogweed), White Clover, Sub-Clover		1.2 - 1.8L + Conquest Glyphosate 450 Herbicide (Glyphosate) and Surfactant at recommended label rates	
	Amaranth, Apple of Peru, Mexican Poppy, Annual Ground Cherry, Bladder Ketmia, Fat Hen, Melons, Native Rosella, Noogoora Burr, Potato Weed, Cow Vine, Yellow Vine, Rapeseed	NSW, ACT, Qld only	1.8 - 2.7L + Conquest Glyphosate 450 Herbicide (Glyphosate) and Surfactant at recommended label rates	
Pastures: Conservation Tillage - Direct Drilling,	Charlock, Mustards, Shepherd's Purse, Saffron, Slender, Spear and Variegated Thistles Turnip Weed, Wild Radish, Wild Turnip	All States	1.1-3.3L	Apply to actively growing young weeds before sowing. Observe plant back periods given in the table on this leaflet.
Surface Sowing or Fallow Maintenance	Clover Sorrel		2.3L plus 280mL-400mL dicamba 500	Apply to actively growing plants in autumn. Do not sow pasture seed for at least 30 days after application.

2. FIELD CROPS

SITUATION & CROP	WEEDS	STATE	RATE/ha	CRITICAL COMMENTS
Wheat	Refer Weed Table	NSW, ACT, SA, Vic, Qld, Tas Only	675mL-2.9L Refer to weed table for Specific rates in each state	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, SA only) Apply from tillering to boot stage (Vic only) Apply from mid-tillering to before boot stage (Qld only) Apply at 5 leaf to fully tillered (Tas only)
Barley				
Cereal Rye, Triticale				
Oats		NSW, ACT, SA Vic, Qld only		
Cereals: Wheat, Oats, Barley	Cape Tulip	WA only	1.4 L - 2.6L	Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33) Apply after the 6 leaf stage (Z 16) for Cranbrook, Jacup, Aroona and Spear Wheat and Mortlock Oats to avoid possible damage. DO NOT spray if lucerne is present. WEED STAGE: 10-15cm. Docks should be sprayed before 5 leaf stage. Cape Tulip - low rate for cornils only
	Dock, Saffron Thistle		2.3L	
	Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Wild Radish		1.6L	
	Wild Turnip		1.4L	
	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip		375mL/ha plus 500mL/ha Diuron 500	
Wheat, Barley	Wild Radish	Vic, SA NSW, ACT only	165mL/ha plus 850g/ha Methabenzthiazuron	Spray 2-6 weeks after sowing and not later. Do not use on crops undersown with lucerne.
Fallow, Stubble Spray prior to Direct Drilling or Sowing a) Winter Cereals b) Winter Cereals and Maize, Sweet Corn, Peanuts Maize, Sweetcorn	Refer Weed Table	Vic only	465mL - 2.9L	Observe plant back periods given in the table in this booklet. Can be mixed with Chlorsulfuron, Paraquat or Paraquat/Diquat where grasses are present. For skeleton weed spraying should only be done 6-8weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
		NSW, ACT only	1.2 - 3.5L	
		Qld only		
Maize, Sweetcorn		NSW, ACT, SA Only	1.2L - 2.3L	Apply when crop is 10 - 20cm high and secondary roots are developing for an over the top spray. When crop is between 20cm high and just before tasselling, spray with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves
		Qld only		Apply when crop is 10 - 30 cm high and secondary roots have developed. Use drop nozzles and direct spray when crop exceeds 30cm in height and before tasselling.
		Tas only		Apply when the crop is 15 - 30cm high. Do not spray if the crop is showing signs of stress. Some leaf twisting may occur following application - crop recovers quickly.

2. FIELD CROPS (continued)

SITUATION & CROP	WEEDS	STATE	RATE/ha	CRITICAL COMMENTS
Sorghum		NSW, ACT, SA Vic only	1.2L – 2.3L	Apply preferably when crop is at 3 - 6 fully expanded leaf stage but can be sprayed from 2 – 8 leaf stage. From 6 leaf stage onwards to within 2 weeks of flowering, crop can be sprayed with dropped nozzles to avoid chemical being sprayed into the whorl and on upper leaves.
		Qld only (except Central Qld)		Apply when crop has 4 to 8 fully expanded leaves and secondary roots have developed.
Millet		NSW, ACT, SA Vic only		Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers.
		Qld only		Do not use on Panarama or Panicum.
Maize, Sweetcorn, Saccaline, Broom Millet, Millet	Cape Tulip, Dock, Saffron Thistle, Indian Hedge Mustard, London Rocket, Lupin,	WA only	2.3L	Spray when crop is 10 - 30 cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds
Grain Sorghum	Rapistrum, Radish, Wild Turnip			Apply when crop is 12cm high. DO NOT apply between tassel and dough stage. Avoid spraying when in flower.
Sugar Cane	Bindy Eye (Star Burr), Blue Top, Cobblers Pegs, Fleabanes, Jute, Leucas, Needle Burr, Spear Thistle, Water Primrose, Ipomea Vines, Convolvulus Vines	Qld only	3.6 – 7.3L	Add 170ml of a 600g/L non-Ionic Surfactant to 100L of spray mixture. Agitate well. DO NOT USE ON Q63, Q67, Q80 OR Q96 VARIETIES
	Chinese Mint, Blue Snakeweed		7.3L	
Peanuts	Broadleaf Weeds, except Noogoora Burr. Grasses, except Mossman Burr	Qld only	3.6L or 7.5L	LOWER RATE Apply as a BAND SPRAY as soon as possible after planting in a 55 cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.
Harvest Aid or Salvage Spray - Winter Cereals	Desiccate Broadleaf Weeds	Qld, NSW, ACT only	2.5 – 3.4L	Apply after dough stage
- Maize & Sorghum	Refer Weed Table			

3. PASTURES, NON AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL LAWNS

SITUATION & CROP	WEEDS	STATE	RATE/ha	CRITICAL COMMENTS
Pastures & Non Agricultural	Refer Weed Table	NSW, ACT, Qld, SA, Tas only	1.1 – 3.5L	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred
	Amsinckia, Docks, Bindweed, Caltrop, Flatweed, Spear Thistle, Capeweed, Doublegee, Saffron Thistle, Mustard, Wild Radish, Wild Turnip, Annual Thistles, Paterson's Curse	WA only	2.3L	For pastures not containing legumes. Only seedling Docks, Spear Thistle and Saffron Thistle will be controlled.
	Afghan Melons		3.3L/ha + 1% crop oil	Spray when plants are actively growing preferably before flowering or vining.
	Paddy Melons		1.6 – 2.3L	
	Prickly Saltwort (Roly Poly)		3.3L	Spray when plants are small
	Stinkwort		3.3 – 6.6L plus surfactant	Best results are obtained when plants are small. Use high rate on larger plants
	Dove weed		6.6L	Spray after good emergence of seedlings
Pastures, Rights of Way & Industrial	Boxthorn, Boneseed, Hawthorn	Vic, SA only	165mL/10L water	Spot spraying. For Boneseed only, thoroughly wet plants or seedlings
	Groundsel	NSW, ACT, Qld, SA only	2.0L/15L water	MISTING: Lightly wet plants
			600mL/100L water	HIGH VOLUME: Thoroughly wet plants
			500mL/15L water	CUT STUMP: Swab the cut stump within one hour of cutting. Apply by a pouring can or knapsack spray.
			6.0 - 9.1L	AERIAL APPLICATION: Spray when Groundsel is actively growing
	Lantana		600mL/100L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet summer (March to May). Defoliation should occur but respraying of new growth will be necessary in the following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.
	Mother of Millions	NSW, ACT only	825ml/100L water	Hand gun and Knapsack only. A thorough coverage of leaves and plantlets is necessary. Use Conquest Wetter 1000 at the rate of 1.0mL or surfactant per 1L of mixture.
	Noogoora Burr, Weir Vine (Ipomea)	Qld only	330mL/100L water	In all cases apply to young, actively growing weeds, ensuring thorough Coverage *spray rosette stage # Repeat spraying if necessary
	Annual & Perennial Pigweed, Artichoke Thistle, Bathurst Burr, Billygoat Weed, Blue Snake Weed, Burr Medic, Clockweed*, Fleabane, Galvanised Burr, Hemlock, Hoary Cress, *Kyalinga Weed (Whisker Grass), Knobweed, Milky Cotton Bushes, Parthenium Weed, Paterson's Curse, Saffron Thistle, Star Burr, Thornapple, Variegated Thistle*		600mL/100L water	
	Rubber Vine		330mL/10L water	

3. FASTURES, NON AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL LAWNS (continued)

SITUATION & CROP	WEEDS	STATE	RATE/ha	CRITICAL COMMENTS
Pastures - Spray/Graze Techniques				PRECAUTION. An increased quantity of poisonous plants may be eaten by stock using Spray-Graze eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse.
	Amsinckia, Thistles, Capeweed, Doublegee, Mustard, Paterson's Curse, Wild Turnip, Wild Radish, Docks, Geranium, Erodium	SA only	1.1L	Apply from 6 weeks after opening rains in autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing. Then return to normal stocking levels. Use high stocking rates following spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control.
	Annual Thistles, Capeweed, Doublegee, Mustards, Paterson's Curse, Turnip, Saffron Thistle, Spear Thistle	Tas, Vic only		
	Amsinckia, Docks (seedlings only), Capeweed, Doublegee, Mustard, Wild Radish, Wild Turnip, Paterson's Curse, Annual Thistles	WA only	1.3L	
	Spear Thistle, Saffron Thistle		2.5L	Apply to Saffron Thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
	Melons		3.3L	Heavy stocking on young plants sprayed with 750mL/ha provides effective control
	Docks	Vic only	2.3L	Apply in September only and follow other recommendations above
	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT only	600mL-2.3L	Spray actively growing 6 - 8 week old weeds. Introduce stock 7 - 10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. Other legumes may be affected.
Lawns	Refer Weed Table	WA & Qld only	3.4 - 6.8mL/1L water	Wet foliage thoroughly

4. SPOT SPRAYING

SITUATION & CROP	WEEDS	STATE	MIXING RATE/CRITICAL COMMENTS
High Volume Spraying	Refer To Weed Table	All States	Add 1/10th of rate in weed table to 150 litres of water. Each 150 litres of mix will cover 1000m ² (1/10th/Ha). Eg. If rate in weed table is 1.5L use 150mL/150L water
Knapsack Application			Add 1/100th of rate in weed table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100th/Ha). Eg. If rate in weed table is 1.5L use 15mL/10L water

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CROP HARVESTING: NOT REQUIRED WHEN USED AS DIRECTED

IN TASMANIA THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WEED TABLE

The rates listed in the Weed Table below are for spot spraying use in crop or pasture, or for use where **weeds only** are present and **no** crop or pasture is involved.

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, (other than spot spraying) use only the rates given for the particular crop or situation indicated under the Directions For Use.

WEEDS	APPLICATION RATE PER HECTARE							CRITICAL COMMENTS
	CROP						PASTURE	
	VIC	NSW, ACT	SA	QLD	TAS	WA	NSW, ACT, SA, QLD, TAS ONLY	
Amaranthus spp.	-	1.1-2.3L	-	1.8L	-	-	-	Spray young plants.
Apple of Peru	-	1.1-2.3L	-	1.8L	-	-	-	Spray young plants. Susceptible when young.
Bathurst Burr	-	1.6-2.3L	-	1.8L	-	-	1.6-2.3L Not SA	Spray seedlings only.
Blackberry Nightshade	-	1.1-2.3L	-	1.8L	-	-	-	
California Burr	-	1.6-2.3L	-	1.8L	-	-	1.6-2.3L not SA	Spray seedlings only.
Cape Tulip	-	-	-	-	-	1.4-2.6L	-	Low rate for cormils only*.
Capeweed	2.9L	-	3.5L	-	2.9L	-	3.4-5.8L	Spray seedlings to rosette stage.
Caltrop	-	1.6-3.5L	-	1.8L	-	-	-	Moderately susceptible
Charlock	1.1-1.6L	1.1-2.3L	1.1L	-	2.9L	-	1.6-2.3L	Spray at rosette stage
Clover	-	2.5L	-	-	-	-	-	
Common Ice Plant	-	-	2.3L	-	-	-	-	
Docks	2.9L	-	2.9L	2.9L	2.9L	2.3L	6.6L SA only	Spray at multiple leaf stage - effective only on seedlings.
Fat Hen	-	1.2-3.5L	-	1.8L	2.9L	-	-	Spray pre-flowering.
Fumitory - red	-	-	3.5L	-	-	-	-	
Fumitory - white	1.6L	-	1.1L	-	-	-	-	Spray at multiple leaf stage.
Hexham Scent or Melilotus	2.9L	-	2.3L	2.9L	-	-	2.3-3.4L	Spray multiple leaf stage before seeding.
Hoary Cress	1.8-2.9L	2.3-3.4L	2.9L	2.9L	-	-	3-3.4L	Spray rosettes at pre-flowering.
Hogweed/ Wireweed	2.9L	-	-	2.9L	-	-	-	Spray at multiple leaf stage (Vic) Spray at seedling and young plant stage (Qld).
Horehound	-	-	2.9L	-	-	-	4.6-6.6L SA only	Spray seedlings.
Khaki Weed	-	-	-	-	-	-	2.3-4.5L not SA	Spray seedlings only
Lincoln Weed	-	-	3.4L	-	-	-	-	Spray early rosettes
London Rocket	-	-	-	-	-	1.6L	-	
Lupins	-	1.6-3.4L	-	-	-	-	-	
Mexican Poppy	-	-	-	2.9L	-	-	-	Spray seedlings - plants become more resistant with age.
Mintweed	-	2.3L	-	1.8L	-	-	-	Spray seedlings - resistant in later stages.
Mustards	465mL-1.1L	1.1-2.3L	1.1 2.9L	1.8L	-	1.6L	1.1-2.3L	Spray at 2-4 leaf up to rosette stage.
New Zealand Spinach	-	2.3-3.4L	-	-	-	-	-	

WEEDS	APPLICATION RATE PER HECTARE							CRITICAL COMMENTS
	CROP						PASTURE	
	VIC	NSW, ACT	SA	QLD	TAS	WA	NSW, ACT, SA, QLD, TAS ONLY	
Noogoora Burr	-	1.6-2.3L	-	1.8L	-	-	1.6-2.3L not SA	Spray seedlings only
Paterson's Curse	-	2.3-3.4	-	2.9L	-	2.6L	3.4-4.6L	Spray rosettes or before plants have 10 leaves. Later stages harder to kill.
Potato Weed	-	1.1-2.3L	-	1.8L	-	-	-	
Rapeseed	-	1.6-3.4L	-	-	-	-	-	
Rough Poppy	-	2.3L	-	-	-	-	-	
Safflower	-	1.1-2.8L	-	-	-	-	-	
Shepherds Purse	-	2.3-3.4L	-	-	2.9L	-	1.6-2.3L	Spray young rosettes
Skeleton Weed	2.9L	2.3-3.4L	2.9L	-	-	-	3-4.6L	Spray rosettes before aerial growth commences.
Sorrel	2.9L	3.4L	2.9L	-	-	-	-	Only moderately susceptible
Speedwell - Ivy leaf	-	-	2.3L	-	-	-	-	
Spiny Emex	-	-	-	2.9L	-	-	-	Only young plants are susceptible.
Stinkwort	-	1.6-2.8L	-	-	-	-	-	
Storksbill/Erodium	-	-	-	-	2.9L	-	3.3-6.6	Spray seedlings to young rosettes.
Sunflower (seedlings)	2.9L	1.1-2.8L	-	1.8L	-	-	-	
Thistles								
- Californian	-	-	-	-	1.2L	-	6.6-7.7L	Repeated applications may be necessary (NSW, Tas only).
- Saffron	2.3L	1.1-2.8L	2.9L	2.9L	2.0L	2.3L	2.3-3.4L	Low rate only sufficient to control weeds in crops at rosette stage when sprayed early.
- Slender/Shore	-	1.6-2.8L	-	-	2.9L	-	2.3L	Suppression only.
- Soldier	2.9L	-	-	-	-	-	2.3-3.3L not NSW, Tas	Spray young rosettes
- Spear	1.1L	-	-	-	2.9L	-	2.3-3.3L	Spray young rosettes
- Star	-	-	-	-	-	-	3.3-6.6L SA only	Use higher rate as flower stalk appears.
- Variegated	-	1.1-3.5L	-	1.8L	2.9L	-	2.3-3.4L	Spray at rosette stage.
Thornapple	-	1.6-2.3L	-	-	-	-	3.5-5.0L not SA	Spray seedlings only.
Turnip Weed/Rapistrum	-	1.1-2.3L	-	1.1L	-	1.6L	1.1-2.3L	
Vetches/Tares	2.9L	-	2.3L	-	-	-	-	Spray at multiple leaf stage.
Wards Weed	-	-	2.3L	-	-	-	-	
Wild Cabbage	2.9L	-	-	-	-	-	-	Spray multiple leaves.
Wild Poppy	1.1L	-	-	-	-	-	2.3-3.4L	Spray rosettes.
Wild Radish	2.9L	3.5L	2.9L	1.8L	2.9L	1.6L	1.6-2.3L	Spray up to young rosette stage.
Wild Turnip	457mL-1.1l	1.1-2.3l	660mL	-	2.9L	1.4L	1.1-2.3L	Spray 2-4 leaf up to rosette stage.

PLANT BACK DAYS FOR CONQUEST AMINE 300

CROP	RATES		
	Up to 1.1 L/ha	1.1 - 2.3 L/ha	2.3 - 3.4 L/ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Bean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

IMPORTANT - WHEN APPLIED TO DRY SOILS, AT LEAST 15 MM (1/2 INCH) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale.

In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall of at least 15 mm

@ In Central Queensland and when using 1.6L/ha or less of Conquest Amine 300 Selective Herbicide, the plant back period for Sorghum and Sunflower is 1 day irrespective of rainfall.

+ In WA the plant back period for Lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Conquest Amine 300 Selective Herbicide is a water soluble liquid product with non-selective herbicidal activity against broadleaf weeds.

Conquest Amine 300 Selective Herbicide will control emerged weeds only, and provides no residual control although certain plant back periods should be observed.

Conquest Amine 300 Selective Herbicide is absorbed by plant foliage and accumulates to toxic levels in the regions of growth and reproduction, upsetting the ability of plants to balance the synthesis and use of nutrients. Visible effects are a gradual yellowing and wilting of the plants which advances to complete browning of above ground growth and deterioration of root systems. Effects may not be apparent for 7 -10 days or even up to 21 days under cold or cloudy conditions.

DO NOT treat weeds under poor growing or dormant conditions such as occur in drought, water-logging disease, insect damage, following frost, weeds heavily covered with dust or silt. Reduced results may also occur if weeds are under stress from previous herbicide application. Rainfall occurring up to 6 hours after application may reduce effectiveness. DO NOT spray if strong winds prevail.

Resistant Weeds Warning

GROUP

HERBICIDE

Conquest Amine 300 Selective Herbicide is a member of the Phenoxy group of herbicides. It has the disruptors of plant cell growth mode of action. For weed resistance management Conquest Amine 300 Selective Herbicide is a group I herbicide.

Some naturally-occurring weed biotypes resistant to Conquest Amine 300 Selective Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individual can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Amine 300 Selective Herbicide or any other group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Agrochemicals Pty. Ltd. accepts no liability for any losses that may result from the failure of Conquest Amine 300 Selective Herbicide to control resistant weeds.

Crop Establishment

Conquest Amine 300 Selective Herbicide is recommended as a herbicide additive to glyphosate for control of emerged weeds prior to crop establishment. When Conquest Amine 300 Selective Herbicide is applied prior to crop establishment, certain Plant Back Periods should be observed (see section on Plant Back Periods) to ensure that the herbicide has degraded sufficiently to allow safe sowing of the intended crop. This process is largely influenced by moisture, temperature and certain soil characteristics and may be delayed particularly when conditions are cold and dry. Refer to the Plant Back Period table for specific information. In seasons of heavy weed growth, or where the following conditions apply, it may be necessary to further delay sowing until a suitable seedbed can be formed. Conditions which can delay crop germination and seedling development include:

- Heavy green or decaying weed growth incorporated into the soil;
- Soil compaction or crusting;
- Cold and wet soils;
- Deep seeding;
- Prior use of residual or pre-emergent herbicides.

To minimise these effects it is suggested that;

- Weed bulk is reduced by grazing and cultivating to leave trash on the surface to dry out;
- A friable seedbed be produced by cultivation, where necessary;
- The use of pre-emergent herbicides to be avoided if they might contribute to reduced germination.
- A correct seeding depth be used.

The preferred alternative is to spray early to control any weeds in their less advanced stages and ensure the seedbed is in a suitable condition for early sowing when soil temperatures are not excessively cold.

Application

Boom Equipment

Application of Conquest Amine 300 Selective or Conquest Glyphosate 450 Herbicide (Glyphosate) mixtures in spray volumes of 25-100L/Ha is recommended.

When sulfonylureas are included in the spray mixture a minimum volume of 30 L/Ha is recommended. When Simazine is included in the mixture a minimum spray volume of 100L/Ha is recommended. Flat fan nozzle equipment is recommended using pressures in the range 240-280 kPa. Boom height must be set to ensure double overlap of nozzle patterns at the top of the weed canopy.

Aerial Equipment

Application of Conquest Amine 300 Selective or Conquest Glyphosate 450 Herbicide (Glyphosate) mixtures using either Micronair or boom equipment should occur in a minimum spray volume of 15L/Ha. Droplets with an average size of 250-350 micron diameter are recommended. Swath width should be 15-17 metres.

Application under hot conditions: High temperature and/or low relative humidity cause excessive evaporation of spray droplets, which may reduce results. When temperatures reach 25°C increase water volume to 30L/Ha, and increase droplet size to at least 300-micron diameter.

DO NOT apply by aircraft when temperature is above 35°C
DO NOT use in intensive horticultural cropping areas,
Thoroughly wash aircraft, especially landing gear after each day of spraying to remove herbicide residues.

Equipment Maintenance

Spray solutions of Conquest Amine 300 Selective Herbicide and glyphosate should be mixed, stored and applied only in stainless steel, aluminium, brass, copper, fibreglass, or plastic lined containers. Do not mix, store or apply spray solutions in galvanised steel or unlined steel (except stainless steel) containers or spray tanks. Conquest Amine 300 Selective Herbicide/glyphosate spray solutions may react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture that can flash or explode if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Thoroughly clean all equipment after use either by using hot soapy water or 1% solution of ammonia followed by several clean water rinses or use of a Tank & Equipment Cleaner.

If using Conquest Chlorsulfuron 750 WG Herbicide or Conquest Metsulfuron 600 WG Herbicide (sulfonylurea herbicides) follow decontamination procedures detailed on those product labels.

Compatibility

Conquest Amine 300 Selective Herbicide can be tank mixed with the products Conquest Glyphosate 450 Herbicide (Glyphosate) Conquest Chlorsulfuron 750 WG Herbicide (Chlorsulfuron), or Conquest Metsulfuron 600 WG Herbicide (Metsulfuron Methyl), Chlorpyrifos as well as Dicamba aqueous concentrates, Simazine and Atrazine Flowables, Paraquat/Diquat, Dimethoate and Omethoate.

Surfactant Addition

DO NOT add a surfactant except in conservation tillage where the product is to be tanked-mixed with products Conquest Glyphosate 450 Herbicide (Glyphosate) or Glyphosate 360. In this situation always add Conquest Wetter 1000 non-ionic surfactant in accordance with label directions on the glyphosate product.

DO NOT mix with spraying oils, or any other materials or agricultural chemicals except as directed on this label.

Tank Mixtures

The Conquest Amine 300 Selective Herbicide directions for use on this label are designed to be used as a tank mixture with Conquest products product Conquest Glyphosate 450 Herbicide (Glyphosate). However as shown in the compatibility and surfactant addition sections of this label, it is possible to extend/improve weed control to include other foliage applied and/or residual herbicides and adjuvants.

A mixture of Conquest Amine 300 Selective Herbicide and Glyphosate may be tank mixed with the following herbicides, insecticides and adjuvants where recommended in the Directions For Use tables. Read and follow all label directions, restraints and plant back periods and safety directions for the tank mix products.

Dicamba 500g/L aqueous concentration.

For improved control of Sow Thistle. Observe any regional use restrictions.

Conquest Chlorsulfuron 750 WG Herbicide (Chlorsulfuron)

Will provide control for a wide range of broadleaf weeds and grasses.

Conquest Metsulfuron 600 WG Herbicide (Metsulfuron Methyl)

For improved knockdown control of Yellow Burrweed, (Amsinckia), Volunteer Chickpeas, Chickweed, Common Sowthistle, Cut-leaf Mignonette, Deadnettle, Faba Beans, Mallee Catchfly, Soursob, Stagger Weed, Wild Garlic. Conquest Metsulfuron 600 WG Herbicide does not provide residual in-crop weed control.

Mixing Instructions

Conquest Amine 300 Selective Herbicide mixes readily with water. Ensure the spray tank is free of any residue of previous spray materials.

1. Fill the spray tank with clean water to one half of the required amount and start agitation. Do not use mechanical agitators as these may cause excessive foaming when herbicides are added.
2. Add recommended herbicide additive/insecticide to the spray tank and mix thoroughly.
3. Add Conquest Amine 300 Selective Herbicide and mix thoroughly
4. Add Conquest Glyphosate 450 Herbicide (Glyphosate).
5. When Conquest Wetter 1000 non-ionic surfactant is used, add near the end of the filling process to minimise foaming.
6. Always maintain adequate agitation during application and use the tank mix promptly.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT use unless windspeed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site
DO NOT apply with smaller than coarse to very coarse spray droplets according to the ASAE S572 definition for standard nozzles

Drift Warning: Direct spray contact, or even slight drift may cause severe injury or destruction of any growing crop or other desirable plant including trees and native vegetation.

DO NOT use Conquest Amine 300 Selective Herbicide by Conquest when breeze is blowing toward nearby desirable plants.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Equipment settings which produce fine droplets (150 microns or less), winds over 8km/hr, "still air" and hot dry days all contribute to drift. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tomatoes, flowers, vegetables, vines, fruit trees, legume crops, pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, Belahs).

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions For Use.

PROTECTION OF WILDLIFE, FISH CRUSTACEAN AND ENVIRONMENT:

DO NOT contaminate dams, rivers or streams with this product or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry, well-ventilated area away from children, animals and food, feedstuffs, seed and fertilisers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers : Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Tel: 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

**FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL
1 800 033 111
24 HOURS AUSTRALIA WIDE.**

Conquest Agrochemicals Pty Ltd.

ABN 84 098 814 932
Suite 9, 41 Walters Drive
Herdsman Business Park
Osborne Park WA 6017

Telephone (08) 9244 1268
Facsimile: (08) 9446 4696

Approval No: 55701/0214