

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Monepantel is a member of the class of anthelmintics called the Amino-Acetonitrile Derivatives (AADs) while Abamectin is a member of the Macrocytic Lactone (ML) class of anthelmintics. ZOLVIX PLUS is effective for the treatment and control of sensitive strains of nematodes (roundworms), including nematodes that have single, dual or triple resistance to macrocytic lactones (including moxidectin resistant strains), benzimidazoles (white drenches), imidazothiazoles (levamisole and morantel (clear drenches) and closantel (*Haemonchus contortus* only), or their combinations in sheep.

EFFICACY SPECTRUM

Barber's pole worm *Haemonchus contortus***^{*}, Small brown stomach worm *Ostertagia (Teladorsagia) spp.***^{*}, *Ostertagia (Teladorsagia) circumcincta**, *Ostertagia (Teladorsagia) trifurcata**, *Ostertagia (Teladorsagia) davitiani**, *Ostertagia (Teladorsagia) spp.***^{*}, Black scour worm *Trichostrongylus colubriformis**, *Trichostrongylus vitrinus**, *Trichostrongylus rugatus**, Stomach hair worm *Trichostrongylus axei***^{*}, Thin necked intestinal worm *Nematodirus filicollis**, *Nematodirus spathiger**, Small intestinal worm *Cooperia curticei**, *Cooperia oncophora**, Large mouthed bowel worm *Chabertia ovina**, and Large bowel worm *Oesophagostomum venulosum**.

* Includes adult and immature L4 stage

** Includes adult, immature L4 stage and inhibited L4 stage

Elanco, Zolvix and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Broad Spectrum Oral Anthelmintic for Sheep

ACTIVE CONSTITUENTS:
25 mg/mL Monepantel
and 2 mg/mL Abamectin

Elanco Australasia Pty Ltd
112 Wharf Road, West Ryde
NSW 2114, Australia.



YL090016ALA 311250

10 L



25 kg x 4000

READ THE ENCLOSED LEAFLET FOR FULL INSTRUCTIONS

DIRECTIONS FOR USE

Restraints: DO NOT USE in sheep less than 10 kg of bodyweight. DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Contraindications: This product is contraindicated for use in lambs under 6 weeks of age or less than 10 kg body weight.

Dosage and Administration: Use remaining product within 12 months of opening the container.

ZOLVIX™ PLUS is a ready-to-use oral solution.

Check dose rates and drench gun before treatment commences. The dose for sheep and lambs is 0.5 mL/5 kg (1 mL/10 kg), equivalent to 2.5 mg monepantel/kg liveweight and 0.2 mg abamectin/kg liveweight. Drench sheep orally using an Elanco recommended drench gun. Maintain applicator gun carefully to ensure accurate dosage. After use, clean gun by flushing with warm, soapy water. Rinse with cold water.

Body weight (kg)	Dose (mL)	No. Treated/ 10 L
10–15	1.5	6666
16–20	2.0	5000
21–25	2.5	4000
26–30	3.0	3330
31–35	3.5	2850
36–40	4.0	2500
41–50	5.0	2000
51–60	6.0	1660
61–70	7.0	1420
71–80	8.0	1250

Sheep in excess of 80 kg body weight should be dosed at 1 mL/10 kg.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest sheep by liveweight in each group (ewes, wethers, rams, lambs). DO NOT under dose. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. A correct drenching technique is recommended; carefully place the drench gun nozzle in the side of the mouth and over the back of the tongue before delivering the dose.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export. Before using this product, confirm the current ESI from Elanco Animal Health on 1800 226 324 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS: Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When opening the container and using the product, wear elbow length chemical-resistant gloves. If product on skin, immediately, wash area with soap and water. Wash hands after use. After each day's use wash gloves.

FIRST AID: If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET: If additional hazard information is required, refer to the Safety Data Sheet. For a copy phone 1800 226 324.

ENVIRONMENTAL PROTECTION: Protection of Wildlife, Fish, Crustacea and

Environment: Do not contaminate dams, waterways or drains with the product or used

containers. Do not discharge waste liquid into streams.

DISPOSAL: The container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced by may be taken separately. DO NOT burn empty containers or product.

STORAGE: Store below 30°C (Room Temperature). Store in the original container tightly closed. Protect from light. Use the remaining product within 12 months of opening the container.

☎ CUSTOMER INFORMATION Australia: 1800 226 324
or visit www.zolvix.com
APVMA Approval No. 69763/106967

Elanco, Zolvix and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.



Batch No.	
Expiry Date	