

COPPERNATE

INJECTION

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING

FOR ANIMAL TREATMENT ONLY

TROY



COPPERNATE

INJECTION

ACTIVE CONSTITUENT:
60mg/mL COPPER as COPPER GLYCINATE

For the prevention and
treatment of copper deficiency
in cattle

NEW VIAL

250ml



TROY LABORATORIES PTY LIMITED
89 LONG STREET SMITHFIELD NSW 2164 AUSTRALIA
web: www.troylab.com.au email: info@troylab.com.au
Dispose of empty container by wrapping
with paper and putting in garbage.

Store below 25°C (air conditioning)
APVMA Approval No.: 60441/V250ML/1105
Batch: Expiry:

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING

FOR ANIMAL TREATMENT ONLY

TROY



COPPERNATE

INJECTION

ACTIVE CONSTITUENT:
60mg/mL COPPER as COPPER GLYCINATE

For the prevention and
treatment of copper deficiency
in cattle

NEW VIAL

250ml

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months
CALVES: 1mL, S.C., repeated within 3-6 months

WITHHOLDING PERIODS: NIL

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with
eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information
Centre. Phone Australia 131 126; New Zealand 0800 764 766

READ THE ATTACHED LEAFLET BEFORE USING:

DIRECTIONS FOR USE:

Excessive copper is toxic; do not administer
unless copper deficiency has been confirmed.
Swelling may occur at injection site which usually
recedes in a few weeks.
Anaphylactic reactions following administration
may occur in young bovines. Prompt antihistamine
treatment is indicated.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TROY

COPPERNATE

INJECTION

ACTIVE CONSTITUENT:

60mg/mL COPPER as COPPER GLYCINATE

For the prevention and treatment of copper deficiency in cattle

250mL



READ THE ATTACHED LEAFLET BEFORE USING:

DIRECTIONS FOR USE:

Excessive copper is toxic, do not administer unless copper deficiency has been confirmed. Swelling may occur at injection site which usually recedes in a few weeks.

Anaphylactic reactions following administration may occur in young bovines. Prompt antihistamine treatment is indicated.

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months

CALVES: 1mL, S.C., repeated within 3-6 months

WITHOLDING PERIODS: NIL

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. **Phone Australia 131126; New Zealand 0800764766.**



TROY LABORATORIES PTY LIMITED
98 LONG STREET SMITHFIELD NSW 2164 AUSTRALIA

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air conditioning).

APVMA Approval No.: 50441/V250ML/1105

Batch:

Expiry:

LT9041

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TROY

COPPERNATE INJECTION

ACTIVE CONSTITUENT:
60 mg/mL COPPER as COPPER GLYCINATE

FOR THE PREVENTION AND TREATMENT OF COPPER DEFICIENCY IN CATTLE
250mL

DIRECTIONS FOR USE:

Excessive copper is toxic; do not administer unless copper deficiency has been confirmed.

Swelling may occur at injection site which usually recedes in a few weeks. Anaphylactic reactions following administration may occur in young bovines. Prompt antihistamine treatment is indicated.

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months

CALVES: 1mL, S.C., repeated within 3-6 months

CAUTION: AVOID CARCASE DAMAGE

1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean; replace frequently.
4. Use shortest needle possible, certainly not exceeding 15 mm.
5. As far as is possible avoid injection of animals in wet weather or under dusty conditions.
6. This product must be injected only under the skin.
7. If possible inject high on the neck behind the ear.

WITHHOLDING PERIODS: NIL



TROY LABORATORIES PTY. LIMITED

98 LONG STREET SMITHFIELD NSW 2164 AUSTRALIA

www.troylab.com.au email: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C.

APVMA 50441/1105

