

DRAFT:
Front Panel:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



TROY
COPPERNATE INJECTION

ACTIVE CONSTITUENT:
60 mg/mL COPPER as COPPER GLYCINATE

**FOR THE PREVENTION
AND TREATMENT
OF COPPER DEFICIENCY
IN CATTLE**

100mL, 250mL



Ancillary Panels:

READ THE ATTACHED LEAFLET BEFORE USING:

DIRECTIONS FOR USE:

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

Excessive copper is toxic; do not administer unless copper deficiency has been confirmed.

Swelling may occur at injection site which usually recedes in a few weeks. Anaphylactic reactions following administration may occur in young bovines. Prompt antihistamine treatment is indicated.

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months

CALVES: 1mL, S.C., repeated within 3-6 months

WITHHOLDING PERIODS: MEAT: NIL MILK: NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): zero (0) days.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. **Phone Australia 131126**

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761 AUSTRALIA

W: www.troylab.com.au E: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning).

APVMA Approval No.: 50441/50148

Batch:

Expiry:

CDXXXX

DRAFT:
Front Panel:

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TROY
COPPERNATE INJECTION

ACTIVE CONSTITUENT:
60 mg/mL COPPER as COPPER GLYCINATE

FOR THE PREVENTION
AND TREATMENT
OF COPPER DEFICIENCY
IN CATTLE

100mL, 250mL



Ancillary Panels:

READ THE ATTACHED LEAFLET BEFORE USING:

DIRECTIONS FOR USE:

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

Excessive copper is toxic; do not administer unless copper deficiency has been confirmed.

Swelling may occur at injection site which usually recedes in a few weeks.
Anaphylactic reactions following administration may occur in young bovines.
Prompt antihistamine treatment is indicated.

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months

CALVES: 1mL, S.C., repeated within 3-6 months

WITHHOLDING PERIODS: MEAT: NIL MILK: NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): zero (0) days.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. **Phone Australia 131126**

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761 AUSTRALIA

W: www.troylab.com.au E: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C (Air Conditioning).

APVMA Approval No.: 50441/50148

Batch:

Expiry:

CDXXXX

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TROY COPPERNATE INJECTION

ACTIVE CONSTITUENT:
60 mg/mL COPPER as COPPER GLYCINATE

**FOR THE PREVENTION AND TREATMENT OF COPPER DEFICIENCY IN
CATTLE**

100mL, 250mL

DIRECTIONS FOR USE:

Use contents within 24 hours of first broaching of the vial. Discard the unused portion.

Excessive copper is toxic; do not administer unless copper deficiency has been confirmed.

Swelling may occur at injection site which usually recedes in a few weeks. Anaphylactic reactions following administration may occur in young bovines. Prompt antihistamine treatment is indicated.

SHAKE WELL BEFORE USE

CATTLE: 2mL, S.C., repeated within 6-9 months

CALVES: 1mL, S.C., repeated within 3-6 months

CAUTION: AVOID CARCASE DAMAGE

1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean; replace frequently.
4. Use shortest needle possible, certainly not exceeding 15 mm.
5. As far as is possible avoid injection of animals in wet weather or under dusty conditions.
6. This product must be injected only under the skin.
7. If possible inject high on the neck behind the ear.

WITHHOLDING PERIODS: MEAT: NIL MILK: NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): zero (0) days.

TROY LABORATORIES PTY. LTD

31-35 Glendenning Road, Glendenning NSW 2761 AUSTRALIA

W: www.troylab.com.au E: info@troylab.com.au

Dispose of empty container by wrapping with paper and putting in the garbage.

Store below 25°C.

APVMA Approval No.: 50441/50148