

Product Name OZTIK POUR-ON TICK INHIBITOR FOR CATTLE  
APVMA Approval No: 58799/118023



Label Name:	OZTIK POUR-ON TICK INHIBITOR FOR CATTLE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 25 g/L FLUAZURON Solvents: 100 g/L N-(N-DODECYL)-2-PYRROLIDONE 100 g/L N-(N-OCTYL)-2-PYRROLIDONE 415 g/L N-METHYL-2-PYRROLIDINONE
Claims:	For strategic control of susceptible cattle tick, Boophilus microplus, (including all known acaricide resistant strains) on beef cattle.  READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT
Net Contents:	2.5 L, 5 L, 15 L
Directions for Use:	
Restrains:	DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.
Contraindications:	This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.
Precautions:	
Side Effects:	
Dosage and Administration:	

General Directions:	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION</p> <ul style="list-style-type: none"> <li>• Re-treat when you first see ticks but not before at least 6 weeks after the previous treatment.</li> <li>• Maximum number of treatments per annual tick season: 3.</li> <li>• Apply in two bands, about 7 cm wide, on each side of the spine between the rump and the shoulders with a suitable applicator.</li> <li>• The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.</li> <li>• There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.</li> <li>• Oztik Pour-On Tick inhibitor for Cattle should not be applied to show cattle as a light scurfiness may be visible for some time after treatment</li> <li>• Oztik Pour-On Tick Inhibitor for Cattle is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of Oztik Pour-On Tick Inhibitor for Cattle commences within 3 days of treatment.</li> <li>• Not suitable for cleaning cattle moving onto "gazetted" cattle tick free areas.</li> </ul> <p>Oztik Pour-On Tick Inhibitor for Cattle treated areas – Introduction of new cattle Tick infested cattle introduced onto properties where an Oztik Pour-On Tick Inhibitor for Cattle treatment regime is in place should be treated with Oztik Pour-On Tick Inhibitor for Cattle prior to release.</p> <p>Handling of Treated Animals Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.</p> <p>For local tick control recommendations contact your local department of agriculture.</p> <p>Cleaning instructions After use wash the applicator with detergent and hot water. Dispose of rinsates in a disposal pit away from desirable plants and their roots and watercourses.</p>
Withholding Periods:	<p>MEAT: DO NOT USE less than 42 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.</p> <p>MILK: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.</p>
Trade Advice:	<p>TRADE ADVICE – Export Slaughter Interval (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website(<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>).</p>
Safety Directions:	<p>Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.</p>

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	If additional hazard information is required refer to the Safety Data Sheet. For a copy phone: 1800 023 312.
-------------------------	--

Environmental Statements:	DO NOT contaminate fish ponds, dams or rivers with the product or used containers. The product is very toxic to aquatic invertebrates.
---------------------------	--

Disposal:	Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------	--

Storage:	Store below 30°C (room temperature). Store in original container, tightly closed in a safe place out of direct sunlight.
----------	--

**DOSE VOLUME TABLE**

<b>Animals and Pest</b>	<b>States</b>	<b>Bodyweight (kg)</b>	<b>Dose (mL)</b>	<b>2.5 L pack treats (head)</b>	<b>5 L pack treats (head)</b>	<b>15 L pack treats (head)</b>
Beef cattle after weaning, Cattle tick ( <i>Boophilus microplus</i> )	Qld, NT, WA and NSW only	101 - 150	9	277	555	1666
		151 - 200	12	208	416	1250
		201 - 250	15	166	333	1000
		251 - 300	18	138	277	833
		301 - 350	21	119	238	714
		351 - 400	24	104	208	625
		401 - 450	27	92	185	555
		451 - 500	30	83	166	500
		501 - 550	33	75	151	454
		551 - 600	36	69	138	416
		601 - 650	39	64	128	384

Add 3 mL for each 50 kg above 650 kg.