



TECHNICAL NOTES

Oztik

Pour-On Tick Inhibitor for Cattle



Active Constituents

Fluazuron 25 g/L

Actions

A tick development inhibitor that kills ticks progressively; therefore, it can take two to three weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. Oztik begins to inhibit tick development within three days of treatment and may give adequate control of ticks for prolonged periods, depending on growth of cattle and the effects of lactation.

Indications

For the strategic control of susceptible cattle tick, *Boophilus microplus*, (including all known acaricide resistant strains) on beef cattle.

Contraindications

Do not treat pregnant cows within six weeks of calving to avoid handling stress.

Restraints

DO NOT USE on cattle which are producing, or may in the future produce milk for human consumption.

Withholding Periods

Meat: DO NOT USE less than 42 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

Milk: DO NOT USE on cattle which are producing, or may in future produce milk or milk products for human consumption.



www.jurox.com.au
Customer Service 1800 023 312
® Registered Trademark of Jurox Pty Limited

AUSTRALIA'S
ANIMAL HEALTH COMPANY





TECHNICAL NOTES

Trade Advice

Export Slaughter Interval (ESI)

DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval.

Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website

(www.apvma.gov.au/residues)

Dosage and Administration

GENERAL INSTRUCTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

- Retreat when you first see ticks but not before at least 6 weeks after the previous treatment.
- Maximum number of treatments per annual tick season: 3.
- Apply in two bands, about 7 cm wide, on each side of the spine between the rump and the shoulders with a suitable applicator.
- The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.
- There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.
- Oztik Pour-On Tick Inhibitor for Cattle is a Tick Development Inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of Oztik Pour-On Tick Inhibitor for Cattle commences within 3 days of treatment.
- Not suitable for cleaning cattle moving into "gazetted" cattle tick free areas.

If handling treated animals within 24 hours of application wear recommended protective clothing (see Safety Directions).

Animals and Pest	States	Bodyweight (kg)	Dose (mL)	5 L pack treats (head)
Beef cattle after weaning Cattle tick (<i>Boophilus microplus</i>)	Qld, WA, NT and NSW only	101-150	9	555
		151-200	12	416
		201-250	15	333
		251-300	18	277
		301-350	21	238
		351-400	24	208
		401-450	27	185
		451-500	30	166
		501-550	33	151
		551-600	36	138
		601-650	39	128

Add 3 mL for each 50 kg above 650 kg.





TECHNICAL NOTES



Safety Directions

Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length butyl rubber gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 1126. If swallowed, DO NOT induce vomiting. Give a glass of water. If product in eyes, wash it out immediately with water: hold eyes open, flood with water for at least 15 minutes and see a doctor. If product on skin, immediately wash area with soap and water.

Presentation

Liquid: 2.5 L, 5 L (backpack) , 15 L

Storage

Store below 30°C (room temperature). Store in original container, tightly closed in a safe place out of direct sunlight.

GHS Information

See Safety Data Sheet for GHS information.

Poisons Schedule

S6

Registration Number

APVMA No. 58799

AU - Tech Note - Oztik / Jul 2019 / APPROVED



www.jurox.com.au
Customer Service 1800 023 312
® Registered Trademark of Jurox Pty Limited

AUSTRALIA'S
ANIMAL HEALTH COMPANY

