

209 mm

190 mm

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

Oztik




POUR-ON TICK INHIBITOR FOR CATTLE

ACTIVE CONSTITUENT: 25 g/L FLUAZURON
Solvents: 100 g/L N - (N-Dodecyl)-2-PYRROLIDONE
100 g/L N - (N-Octyl)-2-PYRROLIDONE,
415 g/L N-METHYL-2-PYRROLIDONE

For Strategic Control of susceptible Cattle Tick,
Boophilus microplus, on Beef Cattle.

**READ DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT**

5L



100% Australian owned,
Australian made

Jurox


408546

APVMA File: 58799_43266_5L_front_MPL_V1.pdf
Dimensions: label = 190mm (high) x 209mm (wide)
Scale: 100% on A4
Date: 14.November.07

100% of final size (190 x 209mm)
209 mm

190 mm

Oztik Pour-On Tick Inhibitor for Cattle



DIRECTIONS FOR USE

Restraints

DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

•This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

•Re-treat when you first see ticks but not before at least 6 weeks after the previous treatment.

•Maximum number of treatments per annual tick season:3

•Apply in two bands, about 7 cm wide, on each side of the spine between the rump and the shoulders with a suitable applicator.

•The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.

•There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother.

A reduced length of protection may occur in cows which are suckling calves.

•Oztik Pour-On Tick Inhibitor for Cattle should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.

•Oztik Pour-On Tick Inhibitor for Cattle is a Tick Development

Inhibitor and kills ticks progressively. It can take 2 to 3 weeks for

cattle to be visibly free of ticks. No additional treatment is

necessary during that time, even on heavily infested animals.

The development inhibiting effect of Oztik Pour-On Tick Inhibitor

for Cattle commences within 3 days of treatment.

•Not suitable for cleaning cattle moving into "gazetted" cattle tick free areas.



DOSE VOLUME TABLE

Animals and Pest	States	Bodyweight (kg)	Dose (mL)	5 L pack treats (head)
Beef Cattle after weaning Cattle Tick (<i>Boophilus microplus</i>)	Qld, NT, WA and NSW only	101 - 150	9	555
		151 - 200	12	416
		201 - 250	15	333
		251 - 300	18	277
		301 - 350	21	238
		351 - 400	24	208
		401 - 450	27	185
		451 - 500	30	166
		501 - 550	33	151
		551 - 600	36	138
		601 - 650	39	128

Add 3mL for each 50 kg above 650 kg

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

MILK: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Ltd on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

General Instructions

Oztik Pour-On Tick Inhibitor for Cattle treated areas - introduction of new cattle.

Tick infested cattle introduced onto properties where a Oztik Pour-On Tick Inhibitor for Cattle treatment regime is in place should be treated with Oztik Pour-On Tick Inhibitor for Cattle *prior* to release.

Handling of Treated Animals

Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

Cleaning instructions

After use wash the applicator with detergent and hot water. Dispose of rinsates in a disposal pit away from desirable plants and their roots and watercourses.

SAFETY DIRECTIONS:

Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.* If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Protection of Wildlife, Fish, Crustaceans and Environment

DO NOT contaminate fish ponds, dams or river with the product or used containers. The product is very toxic to aquatic invertebrates.

Material Safety Data Sheet

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone: (1800) 023 312.

Disposal of Containers and Residual Chemical

Triple rinse containers with detergent and hot water before disposal. Dispose of rinsate in a disposal pit away from desirable plants and their roots and watercourses. Destroy empty containers by breaking, crushing or puncturing them. Dispose of containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area bury the containers at a depth of 500 mm or more at an approved disposal site. DO NOT burn empty containers or product.

STORE below 30°C (Room Temperature). Store in original container, tightly closed in a safe place out of direct sunlight.

APVMA Approval Number 58799/5L/1108

(B)

EXP.



Jurox Pty Limited
85 Gardiner Rd
Rutherford NSW 2320
Infoline 1800 023 312



408554

APVMA File: 58799_xxxx_5L_rear_MPL_V1.pdf
Dimensions: label = 190mm (high) x 209mm (wide)
Scale: 100% on A4
Date: 4th November 08

Bottle Front label

**POISON**

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

Oztik Pour-On Tick Inhibitor for Cattle

Active Constituent: 25 g/L FLUAZURON
Solvents: 100 g/L N-(N-DODECYL)-2-PYRROLIDONE
100 g/L N-(N-OCTYL)-2-PYRROLIDONE
415 g/L N-METHYL-2-PYRROLIDONE

For Strategic Control of the Cattle Tick, *Boophilus microplus*,
(including all known acaricide resistant strains) on Beef Cattle.

READ DIRECTIONS FOR USE BEFORE USING THIS PRODUCT

100% Australian owned,
Australian made

2.5L

jurox

Bottle Back label

DIRECTIONS FOR USE**Restraints**

DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

- Retreat when you first see ticks but not before at least 6 weeks after the previous treatment.
- Maximum number of treatments per annual tick season: 3.
- Apply in two bands, about 7 cm wide, on each side of the spine between the rump and the shoulders with a suitable applicator.
- The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.
- There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves.
- Oztik Pour-On Tick Inhibitor for Cattle should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.
- Oztik Pour-On Tick Inhibitor for Cattle is a Tick Development Inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of Oztik Pour-On Tick Inhibitor for Cattle commences within 3 days of treatment.
- Not suitable for cleaning cattle moving into "gazetted" cattle tick free areas.

DOSE VOLUME TABLE

Animals and Pest	States	Bodyweight (kg)	Dose (mL)	2.5L pack treats (head)
<u>Beef cattle</u>	Qld, NT,	101 – 150	9	277
<u>after</u>	WA &	151 – 200	12	208
<u>weaning</u>	NSW	201 – 250	15	166
Cattle Tick	only	251 – 300	18	139
(<i>Boophilus</i>		301 – 350	21	119
<i>microplus</i>)		351 – 400	24	104
		401 – 450	27	92
		451 – 500	30	83
		501 – 550	33	76
		551 – 600	36	69
		601 – 650	39	64

Add 3 mL for each 50 kg above 650 kg

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

MILK: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Ltd on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

General Instructions**Oztik Pour-On Tick Inhibitor for Cattle treated areas - Introduction of new cattle**

Tick infested cattle introduced onto properties where a Oztik Pour-On Tick Inhibitor for Cattle treatment regime is in place should be treated with Oztik Pour-On Tick Inhibitor for Cattle *prior* to release.

Handling of Treated Animals

Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

Cleaning instructions

After use wash the applicator with detergent and hot water. Dispose of the rinsates in a disposal pit away from desirable plants and their roots and watercourses.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126*. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Protection of Wildlife, Fish, Crustaceans and Environment

DO NOT contaminate fish ponds, dams or rivers with the product or used containers. The product is very toxic to aquatic invertebrates.

Material Safety Data Sheet

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone: (1800) 023 312.

Bottle Back label (continued)

Disposal of Containers and Residual Chemical

Triple rinse containers with detergent and hot water before disposal. Dispose of rinsate in a disposal pit away from desirable plants and their roots and watercourses. Destroy empty containers by breaking, crushing or puncturing them. Dispose of containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area bury the containers at a depth of 500 mm or more at an approved disposal site. DO NOT burn empty containers or product.

Jurox Pty Limited
85 Gardiners Rd
Rutherford NSW 2320
Infoline 1800 023 312

Store below 30°C (Room Temperature). Store in original container, tightly closed in a safe place out of direct sunlight.

APVMA Approval No. 58799/2.5L/xx

B:

Exp:

100% of final size (175 x 200mm)

200 mm

175 mm

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**



Oztik

POUR-ON TICK INHIBITOR FOR CATTLE

ACTIVE CONSTITUENT: 25 g/L FLUAZURON

Solvents: 100 g/L N - (N-Dodecyl)-2-PYRROLIDONE


100 g/L N - (N-Octyl)-2-PYRROLIDONE,

415 g/L N-METHYL-2-PYRROLIDONE

For Strategic Control of susceptible Cattle Tick,
Boophilus microplus, on Beef Cattle.

**READ DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT**

15L



100% Australian owned.
Australian made

Jurox

408756


APVMA File: 58799_43266_15L_front_MPL_V1.pdf
Dimensions: label = 175mm (high) x 200mm (wide)
Scale: 100% on A4
Date: 14.November.07

100% of final size (175 x 200mm)

200 mm

175 mm

Oztik Pour-On Tick Inhibitor for Cattle




DIRECTIONS FOR USE

Restraints

DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

- This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.
- Re-treat when you first see ticks but not before at least 6 weeks after the previous treatment.
- Maximum number of treatments per annual tick season: 3
- Apply in two bands, about 7 cm wide, on each side of the spine between the rump and the shoulders with a suitable applicator.
- The weight of cattle should be determined by either scales or a weighband. Dose individuals according to the heaviest animal in the mob. DO NOT underdose.
- There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother.
- A reduced length of protection may occur in cows which are suckling calves.
- Oztik Pour-On Tick Inhibitor for Cattle should not be applied to show cattle as a light scurfiness may be visible for some time after treatment.
- Oztik Pour-On Tick Inhibitor for Cattle is a Tick Development Inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of Oztik Pour-On Tick Inhibitor for Cattle commences within 3 days of treatment.
- Not suitable for cleaning cattle moving into "gazetted" cattle tick free areas.



DOSE VOLUME TABLE

Animals and Pest	States	Bodyweight (kg)	Dose (mL)	15 L pack treats (head)
Beef Cattle after weaning Cattle Tick (<i>Boophilus microplus</i>)	Qld NT, WA and NSW only	101 - 150	9	1666
		151 - 200	12	1250
		201 - 250	15	1000
		251 - 300	18	833
		301 - 350	21	714
		351 - 400	24	625
		401 - 450	27	555
		451 - 500	30	500
		501 - 550	33	454
		551 - 600	36	416
		601 - 650	39	384

Add 3mL for each 50 kg above 650 kg

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

MILK: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Ltd on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

General Instructions

Oztik Pour-On Tick Inhibitor for Cattle treated areas - introduction of new cattle.

Tick infested cattle introduced onto properties where a Oztik Pour-On Tick Inhibitor for Cattle treatment regime is in place should be treated with Oztik Pour-On Tick Inhibitor for Cattle prior to release.

Handling of Treated Animals

Wear protective clothing recommended under Safety Directions, if handling treated animals within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

Cleaning instructions

After use wash the applicator with detergent and hot water. Dispose of rinsates in a disposal pit away from desirable plants and their roots and watercourses.

SAFETY DIRECTIONS:

Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Protection of Wildlife, Fish, Crustaceans and Environment

DO NOT contaminate fish ponds, dams or river with the product or used containers. The product is very toxic to aquatic invertebrates.

Material Safety Data Sheet

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone: (1800) 023 312.

Disposal of Containers and Residual Chemical

Triple rinse containers with detergent and hot water before disposal. Dispose of rinsate in a disposal pit away from desirable plants and their roots and watercourses. Destroy empty containers by breaking, crushing or puncturing them. Dispose of containers at a local authority landfill that does not burn its refuse. If there is no local authority landfill readily available in your area bury the containers at a depth of 500 mm or more at an approved disposal site. DO NOT burn empty containers or product.

STORE below 30°C (Room Temperature). Store in original container, tightly closed in a safe place out of direct sunlight.

APVMA Approval Number 58799/15L/1108

(B)

EXP.



Jurox Pty Ltd
85 Gardiner Road
Rutherford NSW 2320
Infoline 1800 023 312



9 1338094 1000431

408763

APVMA File: 58799_XXXXX_15L_rear_MPL_V1.pdf
Dimensions: label = 175mm (high) x 200mm (wide)
Scale: 100% on A4
Date: 4th November 08