

SF00-SF06-0002

# EPRIGARD

Pour-On for Beef and Dairy Cattle



## Active Content & Action

Eprinomectin 5 g/L. Eprinomectin is part of the Macrocylic Lactone (ML) group of chemicals. Use of ML's induce paralysis and death in the target parasites.

## Use

For the treatment and control of internal and external parasites of beef and dairy cattle and internal parasites of deer. For the treatment and control of gastrointestinal roundworms (including inhibited immature *Ostertagia ostertagi*) and lungworm in cattle. For sucking and biting lice, sarcoptic and choriopic mange mites and buffalo fly in beef and dairy cattle. Aids in control of cattle tick *Rhipicephalus (Boophilus) microplus*. For the treatment and control of gastrointestinal roundworms and lungworm in deer.

## CATTLE

### Gastrointestinal Roundworms:

Parasites	Adults	Immatures	Inhibited L4
<i>Ostertagia ostertagi</i> - small brown stomach worm	•	•	•
<i>O. lyrata</i> - brown stomach worm	•		
<i>Ostertagia</i> spp.	•	•	
<i>Haemonchus placei</i> - barbers pole worm	•	•	
<i>Trichostrongylus axei</i> - stomach hair worm	•	•	
<i>T. colubriformis</i> - intestinal hair worm	•	•	
<i>Trichostrongylus</i> spp.	•	•	
<i>Cooperia</i> spp. - small intestinal worm	•	•	•
<i>C. oncophora</i>	•	•	
<i>C. pectinata</i>	•	•	
<i>C. surnabada</i>	•	•	
<i>Oesophagostomum radiatum</i> - nodule worm	•	•	
<i>Oesophagostomum</i> spp.	•	•	
<i>Nematodirus helvetianus</i> - thin necked intestinal worm	•	•	
<i>Bunostomum phlebotomum</i> - hookworm	•	•	
<i>Trichuris</i> spp. - whip worm	•		

## Lungworm:

Parasites	Adults	Inhibited L4
<i>Dictyocaulus viviparus</i>	•	

## Sucking and Biting Lice:

*Linognathus vituli*

*Hematopinus eurysternus*

*Bovicola (Damalinia) bovis*

*Solenopotes capillatus*

## Mites:

*Chorioptes bovis*

*Sarcoptes scabiei* var. *bovis*

## Buffalo Fly:

*Haematobia irritans exigua*

When used as directed, Eprigard™ Pour-On for Beef and Dairy Cattle also aids in control of: Cattle Tick:

*Boophilus microplus*

(Not recommended for use in strategic tick control programs)

\*The following parasite species are included within each of the relevant genera: *Ostertagia ostertagi*, *O. leptospicularis*, *O. lyrata*, *Cooperia oncophora*, *C. punctata*, *C. surnabada*, *Trichostrongylus axei*, *T. colubriformis*.

## DEER

### Gastrointestinal roundworms:

Parasites	Adults	Inhibited L4
<i>Ostertagia</i> - type spp. *	•	
<i>Trichostrongylus</i> spp.	•	•
<i>Mazamastrongylus</i> spp.	•	
<i>Oesophagostomum</i> spp.	•	

\*Includes *O. mossi/dikmansi*, *O. lyrata*, *O. leptospicularis*, *Spiculoptera spiculoptera*, *Sp. asymmetrica* and *Apteragia quadrispiculata*.

For best results Eprigard™ Pour-On for Beef and Dairy Cattle should be part of a total parasite control program including internal and external parasites. Resistance may develop to any drench. Consult your veterinarian for a complete parasite control program.

## NOTE TO USER:

Rainfall before or after treatment will not affect the efficacy of Eprigard Pour-On.

## SAFETY:

Studies have demonstrated a wide safety margin. Five times the recommended use level had no adverse effect on cattle or deer. Three times the recommended use level had no adverse effect on breeding performance of cows and bulls.



## Doses

400 doses per 20L pack  
(average weight 500 kg cattle)

## Pack Size(s) Available

20L & 5L

## Australian Poisons Schedule

S5

## Presentation

40 drums to a pallet (20L)  
3 backpacks per shipper (5L)

## APVMA Approval No.:

87119



## Period of Protection

Parasites	Prolonged Activity
<i>Ostertagia</i> spp. *	Up to 28 days
<i>Cooperia</i> spp.	Up to 28 days
<i>Dictyocaulus viviparus</i>	28 days
<i>Oesophagostomum radiatum</i>	Up to 28 days
<i>Nematodirus helvetianus</i>	Up to 28 days
<i>Haemonchus placei</i>	21 days
<i>Trichostrongylus</i> spp.	21 days

\*The following parasite species are included within each of the relevant genera: *Ostertagia ostertagi*, *O.leptospicularus*, *O.lyrata*, *Cooperia oncophora*, *C.punctata*, *C.surnabada*, *Trichostrongylus axei*, *T.colubriformis*.

## Dosage and Administration

A representative sample of animals should be weighed with either scales or a weighband. Dose the mob according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, hinds, stags, fawns, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. This product is formulated only for pour-on application to cattle and deer; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead.

The dose rate is 1 mL for each 10kg of liveweight.

## Withholding Periods

Meat – Nil  
Milk – Nil

## Trade Advice

EXPORT SLAUGHTER INTERVAL (ESI): NIL. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA Pty Ltd on 1800 144 884 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))

## Accessories

This product can be used with any pour on applicator designed to apply pour on products to cattle.  
Contact your local reseller for the different options.

