



Product Name: Cyzmic Insecticide
 APVMA Approval No: 87248/128656



Label Name:	Cyzmic Insecticide
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	100 g/L LAMBDA-CYHALOTHRIN
Mode of Action:	GROUP 3A INSECTICIDE
Statement of Claims:	Situations: Domestic, Commercial, Public and Industrial situations. Controls: Cockroaches, Spiders, Fleas, Ants, Flies, Silverfish and Mosquitoes.
Net Contents:	100 mL - 5 L
Restraints:	DO NOT use in farm buildings where grain or crop produce is stored.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	
Trade Advice:	

General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>Mixing</p> <ol style="list-style-type: none"> 1. Half fill the spray tank with water. 2. Shake CYZMIC well before use. 3. Measure the required amount of CYZMIC using the device attached to the cap. 4. Pour the measured dose of CYZMIC into the sprayer. 5. Add the remaining water, close sprayer, shake and pump to correct pressure and apply spray.
Resistance Warning:	<p>Insecticide Resistance Warning GROUP 3A INSECTICIDE</p> <p>For insecticide resistance management CYZMIC Insecticide is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to CYZMIC and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if CYZMIC or other Group 3A insecticides are used repeatedly. The effectiveness of CYZMIC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Adama Australia accepts no liability for any losses that may result from the failure of CYZMIC to control insects. CYZMIC may be subject to specific resistance management strategies. For further information contact your local supplier, Adama representative or local agricultural department agronomist.</p>
Precautions:	<p>PRECAUTIONS</p> <p>Cover all foodstuffs before application. DO NOT allow spray to contact food, food preparation surfaces and utensils. When using in farm buildings do not spray directly onto livestock, their food, water, litter, feed bins or surfaces accessible to stock.</p> <p>BEFORE USE: Remove or cover all exposed foodstuff. Cover all dishes and utensils and surfaces where food is prepared. Remove animals, remove or cover all exposed stock feed, feed and water containers and troughs, places where stock feed is prepared or stored, and milking equipment.</p> <p>AFTER USE: Thoroughly ventilate treated areas. Clean up thoroughly before processing/ serving resumes.</p>
Protections:	<p>PROTECTION OF WILDLIFE FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT discharge the liquid into waterways. Remove or cover fish tanks before use.</p>
Storage and Disposal:	<p>Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>

Safety Directions:	Harmful if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, face-shield and contaminated clothing.
--------------------	--

First Aid Instructions:	If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE

Situation	Pests	Rate/10 L water	Critical Comments
<p>Commercial, domestic, industrial and public buildings including; barracks, factories, farm buildings, food processing/handling establishments, hospitals, hotels and motels, offices, residential homes, restaurants, shops, schools, storerooms, transportation equipment such as aircraft, boats, buses, ships, trains, warehouses and other situations which fall within the above definitions including outdoor or external areas associated with such establishments, on surfaces including; cement - painted, unpainted and rendered, ceramic tiles, painted plasterboard, steel - stainless and enamelled, wood - painted and unpainted.</p>	German Cockroaches (<i>Blattella germanica</i>)	Clean out 50 mL	<p>Apply dilute spray at the rate of 10 L/200 m² or to runoff.</p> <p>Crawling insects Locate possible harbourages and thoroughly apply CYZMIC as a coarse low pressure spray to these areas. Barrier spraying around infestation origin points may increase control.</p> <p>Flying insects Apply as a residual surface treatment. All people and animals should be removed from buildings prior to treatment.</p> <p>Cockroaches, ants and house flies Indoors CYZMIC (50 mL rate only) can deliver residual control of up to 12 months. Outdoors CYZMIC (50 mL rate only) can deliver residual control of up to 3 months. DO NOT apply in areas exposed to rain and/or direct sunlight. Apply in sheltered areas where insects lurk.</p>
		Maintenance 25 mL	
	American Cockroaches (<i>Periplaneta americana</i>)	Clean out 40 mL	
		Maintenance 25 mL	
	Ants, Cat Flea, House Flies	50 mL	
	Mosquitoes, Spiders	40 mL	
Silverfish	25 mL		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION