



Company Name: ADAMA AUSTRALIA PTY LIMITED
Product Name: 3-D Herbicide
APVMA Approval No: 82412/105683



Label Name:	3-D Herbicide
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENTS: 192 g/L BENTAZONE present as the SODIUM SALT 112 g/L MCPA present as the DIMETHYLAMINE SALT 19 g/L DICAMBA present as the DIMETHYLAMINE SALT
Mode of Action:	GROUP C I HERBICIDE
Statement of Claims:	For the control of Mullumbimby Couch and Broadleaf Weeds in Turf as per the Directions for Use
Net Contents:	1 L - 20 L
Restraints:	DO NOT mow turf for at least 3 days before or after spray application. DO NOT apply if rain is expected within 8 hours. DO NOT apply to Lippia, Strawberry Clover, Buffalo or Saltene (Paspalum notatum) lawns.
Directions for Use:	This section contains file attachment. File Name: 3-D Herbicide_Commercial Label_Directions for Use_160202.pdf File Size: 96073 bytes
Other Limitations:	

Withholding Periods:	
-----------------------------	--

Trade Advice:	
----------------------	--

General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>The product mixes readily in hard or soft water. Weeds are killed by absorption through the leaf and root system of susceptible weeds. It is important to apply the recommended spray mixture when the weeds are young and actively growing and soil is moist. Results may take time to become evident. If necessary to respray, allow an interval of 3 weeks. This product when applied at the recommended rates may be used on established turf such as Couch, Blue Couch, Fescue, Carpet, Centipede, Kentucky Blue and Rye Grass.</p>
------------------------------	--

Resistance Warning:	<p>RESISTANT WEEDS WARNING</p> <p>3-D HERBICIDE is a member of the Benzothiadiazoles, Phenoxy and Benzoic Acid groups of herbicides and acts by inhibiting photosynthesis at photosystem II and disrupting plant cell growth. For weed resistance management the product is both a Group C and a Group I herbicide. Some naturally occurring weed biotypes resistant to 3-D Herbicide and other Group C and/or Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group C and/or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, ADAMA Australia Pty Ltd accepts no liability for any losses that may result from failure of 3-D HERBICIDE to control resistant weeds. However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application.</p>
----------------------------	--

Precautions:	
---------------------	--

Protections:	<p>PROTECTION OF LIVESTOCK</p> <p>DO NOT feed grass clippings from treated areas to livestock or poultry.</p> <p>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</p> <p>DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate streams, rivers or waterways with the chemical or used container.</p>
---------------------	--

Storage and Disposal:	<p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a cool well ventilated area away from children, animals, food, feedstuffs, seed and fertiliser. Do not store for prolonged periods in direct sunlight. Triple rinse or preferably pressure rinse containers before disposal. Add rinsings</p>
------------------------------	--

to the spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Safety Directions:

SAFETY DIRECTIONS

Harmful if swallowed. Attacks the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

First Aid Instructions:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:



3-D HERBICIDE
Item 7
Decision ID: DC7-74434865

DIRECTIONS FOR USE

SITUATION	WEEDS CONTROLLED	STATE	RATE	HOW TO APPLY
Lawns and Turf	Bindii (Dwarf Jo-Jo), Chickweed, Clover, Creeping Oxalis, Cudweed, Dandelion, Dock, Flatweeds, Fleabane, Lambs Tongue, Mullumbimby Couch and Sowthistle	All States	<p>SPRAY:</p> <p>Spray 200 mL in 20 L water over 100m².</p> <p>SPOT SPRAY:</p> <p>60 mL in 10 L water and wet weeds thoroughly</p>	<p>Spray annual weeds when young and vigorous and perennial weeds just prior to flowering when soil is moist. Thoroughly wet foliage and avoid excess runoff. Repeat applications 3 to 4 weeks after initial spraying if necessary.</p> <p>To avoid damage to lawns/turf, use only at recommended rates and do not exceed 2 sprayings within 3 to 4 weeks.</p> <p>Note: for effective control of Mullumbimby Couch, spray in warmer months, when weeds are actively growing.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.