



Product Name: VETSENSE IVERMEC Pour-On for Cattle
APVMA Approval No: 93625/139980

Label Name:	VETSENSE IVERMEC Pour-On for Cattle
-------------	-------------------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	5 mg/mL IVERMECTIN
-------------------------	--------------------

Claims:	<p>For the treatment and control of internal and external parasites of cattle.</p> <p>VETSENSE IVERMEC Pour-on for Cattle contains ivermectin, a member of the macrocyclic lactone family of chemicals.</p> <p>It is effective against sensitive strains of the following internal and external parasites: gastrointestinal nematodes <i>Ostertagia ostertagi</i>, <i>Haemonchus placei</i>, <i>Trichostrongylus axei</i>, <i>T. colubriformis</i>, <i>Cooperia oncophora</i>, <i>C. punctata</i>, <i>Oesophagostomum radiatum</i>, <i>O. venulosum</i>, <i>Strongyloides papillosus</i>, <i>Trichuris spp</i>, and <i>Nematodirus spp</i>, lungworms (<i>Dictyocaulus viviparus</i>), sucking and biting lice (<i>Linognathus vituli</i>, <i>Haematopinus eurysternus</i>, <i>Bovicola bovis</i> and <i>Solenopotes capillatus</i>), chorioptic and sarcoptic mange mites, buffalo flies (<i>Haematobia irritans exigua</i>), and cattle ticks (<i>Boophilus microplus</i>).</p> <p>VETSENSE IVERMEC Pour-On at the recommended use level can effectively control infections with <i>Ostertagia spp.</i> (small brown stomach worms) acquired up to 14 days after treatment, <i>Dictyocaulus viviparus</i> (lungworm) acquired up to 28 days after treatment and <i>Haematobia irritans exigua</i> (buffalo fly) for up to 21 days after treatment.</p> <p>Resistance may develop to any chemical.</p>
---------	---

Net Contents:	250 mL 500 mL 1 L 2.5 L 5 L 15 L 20 L
---------------	---

Directions for Use:																											
Restraints:	<p>Do not use in species other than cattle</p> <p>Do not give orally or parenterally - this product is for application to the skin surface only</p> <p>Re-treatment interval: DO NOT RE-TREAT animals for 42 days after last treatment</p>																										
Contraindications:	This product should not be used in dogs, horses or any other pets																										
Precautions:	Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced. Rain falling on cattle in less than 2 hours after dosing may result in reduced efficacy. Reduced efficacy will result if applied to areas of skin with mange scabs or lesions, or with dermatoses or adherent materials, eg. caked mud or manure.																										
Side Effects:																											
Dosage and Administration:	<p>Apply along the topline in a narrow strip extending from the withers to the tailhead. The dose rate is 1mL for each 10kg of liveweight.</p> <table> <thead> <tr> <th>Live weight (kg)</th> <th>Dose Volume (mL)</th> </tr> </thead> <tbody> <tr><td>50-100</td><td>10</td></tr> <tr><td>101-150</td><td>15</td></tr> <tr><td>151-200</td><td>20</td></tr> <tr><td>201-250</td><td>25</td></tr> <tr><td>251-300</td><td>30</td></tr> <tr><td>301-350</td><td>35</td></tr> <tr><td>351-400</td><td>40</td></tr> <tr><td>401-450</td><td>45</td></tr> <tr><td>451-500</td><td>50</td></tr> <tr><td>501-550</td><td>55</td></tr> <tr><td>551-600</td><td>60</td></tr> <tr><td>601-650</td><td>65</td></tr> </tbody> </table> <p>Heavier animals, for example mature bulls, should receive an additional 5 mL for each 50 kg over 650 kg.</p> <p>A representative sample of animals should be weighed before treatment either with scales or a weighband.</p> <p>Dose rate to be based on heaviest cattle in each group. Do not underdose. When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.</p>	Live weight (kg)	Dose Volume (mL)	50-100	10	101-150	15	151-200	20	201-250	25	251-300	30	301-350	35	351-400	40	401-450	45	451-500	50	501-550	55	551-600	60	601-650	65
Live weight (kg)	Dose Volume (mL)																										
50-100	10																										
101-150	15																										
151-200	20																										
201-250	25																										
251-300	30																										
301-350	35																										
351-400	40																										
401-450	45																										
451-500	50																										
501-550	55																										
551-600	60																										
601-650	65																										
General Directions:	<p>The colour of VETSENSE IVERMEC Pour-On for Cattle fades when exposed to light and, depending on the light intensity, fading may occur in less than 30 minutes. This rapid loss of colour does not reflect loss of potency or Ivermectin. However, prolonged exposure to light can also result in gradual decline of Ivermectin potency in formulation.</p> <p>Gives off highly flammable vapour. Keep away from heat, sparks, open flame and other sources of ignition. Keep closed when not in use. Only use in well-ventilated area or outdoors.</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p>																										

Withholding Periods:	MEAT: DO NOT USE less than 42 days before slaughter for human consumption. MILK: Zero (0) days
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Vetsense Pty Ltd on 0245779911 or the APVMA website (apvma.gov.au/residues).
Safety Directions:	Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. If product in eyes, wash out immediately with water. Wash hands after use.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	
Environmental Statements:	ENVIRONMENTAL PROTECTION Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams and other waterways with the chemical or used container.
Disposal:	For containers less than or equal to 1L: Dispose of container by wrapping with paper and putting in garbage. For large containers greater than 1L: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30 °C away from light.