



Ivermec Pour-On for Cattle



Vetsense Ivermec Pour-On is ideal for the treatment and control of internal and external parasites of cattle. Contains Ivermectin, a member of the macrocyclic lactone family of chemicals.

- ✓ Broad parasite coverage
- ✓ Controls internal and external parasites including Cattle Ticks and Buffalo Flies
- ✓ Safe & economical choice
- ✓ Broad safety margin
- ✓ Rainfast

		Ivermec Pour-On
ACTIVE CONSTITUENTS		5 mg/mL IVERMECTIN
SPECIES		Cattle - Beef & Dairy
DOSE RATE		1 mL / 10 kg
ADMINISTRATION		Pour-On
PACK SIZES		250mL, 1L, 2.5L, 5L & 20L
APVMA NO		93625/139980
WITHHOLDING PERIOD		
MEAT		42 days
MILK		Nil
EXPORT SLAUGHTER INTERVAL		42 days



Contact
Vetsense:



PRODUCT INFORMATION

Ivermec Pour-On for Cattle

Active Constituents: 5 mg/mL IVERMECTIN



Vetsense Ivermec Pour-on for Cattle contains Ivermectin, a member of the macrocyclic lactone family of chemicals.

Ivermec is effective against sensitive strains of the following internal and external parasites:

✓ GASTROINTESTINAL NEMATODES

- *Ostertagia ostertagi*
- *Haemonchus placei*
- *Trichostrongylus axei*
- *T. colubriformis*
- *Cooperia oncophora*
- *C. punctata*
- *Oesophagostomum radiatum*
- *O. venulosum*
- *Strongyloides papillosus*
- *Trichuris spp*
- *Nematodirus spp*

✓ LUNGWORMS (*Dictyocaulus viviparus*)

✓ SUCKING AND BITING LICE

- *Linognathus vituli*
- *Haematopinus eurysternus*
- *Bovicola bovis*
- *Solenopotes capillatus*

✓ CHORIOPTIC AND SARCOPTIC MANGE MITES

✓ BUFFALO FLIES (*Haematobia irritans exigua*)

✓ CATTLE TICKS (*Boophilus microplus*)

At the recommended use level, Ivermec can effectively control infections with *Ostertagia spp.* (small brown stomach worms) acquired up to 14 days after treatment, *Dictyocaulus viviparus* (lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (buffalo fly) for up to 21 days after treatment.

RESTRAINTS

Do not use in species other than cattle.

Do not give orally or parenterally - this product is for application to the skin surface only.

Re-treatment interval: DO NOT RE-TREAT animals for 42 days after last treatment.

CONTRAINdications

This product should not be used in dogs, horses or any other pets.

PRECAUTIONS

Cattle should not be treated when hair or hide is wet since reduced efficacy may be experienced. Rain falling on cattle in less than 2 hours after dosing may result in reduced efficacy. Reduced efficacy will result if applied to areas of skin with mange scabs or lesions, or with dermatoses or adherent materials, eg. caked mud or manure.

DOSAGE AND ADMINISTRATION

Apply along the topline in a narrow strip extending from the withers to the tailhead. The dose rate is 1 mL for each 10 kg of liveweight.

Heavier animals, for example mature bulls, should receive an additional 5 mL for each 50 kg over 650 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group. Do not underdose. When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

LIVWEIGHT	DOSE VOLUME	250 mL PACK TREATS	1L PACK TREATS	2.5L PACK TREATS	5L PACK TREATS	20L PACK TREATS
50-100 kg	10 mL	25	100	250	500	2000
101-150 kg	15 mL	16	66	166	333	1333
151-200 kg	20 mL	12	50	125	250	1000
201-250 kg	25 mL	10	40	100	200	800
251-300 kg	30 mL	8	33	83	166	666
301-350 kg	35 mL	7	28	71	142	571
351-400 kg	40 mL	6	25	62	125	500
401-450 kg	45 mL	5	22	55	111	444
451-500 kg	50 mL	5	20	50	100	400
501-550 kg	55 mL	4	18	45	90	363
551-600 kg	60 mL	4	16	41	83	333
601-650 kg	65 mL	3	15	38	76	307

GENERAL DIRECTIONS

The colour of Vetsense Ivermec Pour-On for Cattle fades when exposed to light and, depending on the light intensity, fading may occur in less than 30 minutes. This rapid loss of colour does not reflect a loss of potency in Ivermectin. However, prolonged exposure to light can also result in gradual decline of Ivermectin potency in formulation.

Gives off highly flammable vapour. Keep away from heat, sparks, open flame and other sources of ignition. Only use in well-ventilated area or outdoors.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

MILK: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Vetsense Pty Ltd on 0245779911 or the APVMA website (apvma.gov.au/residues).

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. If product in eyes, wash out immediately with water. Wash hands after use.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

ENVIRONMENTAL PROTECTION

Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams and other waterways with the chemical or used container.

DISPOSAL

For containers less than or equal to 1L:

Dispose of container by wrapping with paper and putting in garbage.

For large containers greater than 1L:

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.


STORAGE

Store below 30 °C (room temperature), closed tight and away from light.

APVMA No.: 93625/139980



1L-Pack Application Guide

STEP 1	STEP 2
<p>Turn bottle upside down to fill the dosing chamber.</p> 	<p>Once chamber is filled, stand bottle upright to level the dosage.</p> 
STEP 3	STEP 4
<p>Tilt bottle to adjust to the correct dosage.</p> 	<p>Once you have the correct dosage, remove the cap.</p> 
STEP 5	STEP 6
<p>Pour out the liquid.</p> 	<p>Apply cap to reseal the bottle.</p> 