



Triclamec ULTRA Pour-On



Vetsense Triclamec ULTRA Pour-On is ideal for the treatment and control of Roundworms, all 3 stages of Liver Fluke and external parasites of beef cattle.

Contains Abamectin, a member of the macrocyclic lactones family of chemicals.

- ✓ Dual actives for improved efficacy
- ✓ Triclabendazole treats against early-immature, immature and mature stages of Liver Fluke
- ✓ Aids in the control of Cattle Tick, Lice and Buffalo Fly
- ✓ Broad control against immature and adult worms

Triclamec Pour-On

ACTIVE CONSTITUENTS	300 mg/mL TRICLABENDAZOLE, 5 mg/mL ABAMECTIN
SPECIES	Beef Cattle
DOSE RATE	1 mL / 10 kg
ADMINISTRATION	Pour-On
PACK SIZES	1L, 2.5L, 5L & 10L
APVMA NO	94653/143094
WITHHOLDING PERIOD	
MEAT	49 days
MILK	Do not use
EXPORT SLAUGHTER INTERVAL	140 days



Contact
Vetsense:



PRODUCT INFORMATION

Triclamec ULTRA Pour-On

Active Constituents: 300 mg/mL TRICLABENDAZOLE, 5 mg/mL ABAMECTIN



Vetsense Triclamec Ultra contains Abamectin, a member of the macrocyclic lactones family of chemicals.

Triclamec is effective against sensitive strains of internal and external parasites. It also contains Triclabendazole, a member of the benzimidazoles family of chemicals, effective against liver fluke.

INDICATIONS

For the treatment and control of adult and immature stages of:

✓ GASTRO-INTESTINAL ROUNDWORMS

- Barber's Pole Worm (*Haemonchus placei*)
- Small Brown Stomach Worm (*Ostertagia ostertagi*)
- Small Intestinal Worm (*Cooperia oncophora*, *Cooperia punctata*)
- Stomach Hair Worm (*Trichostrongylus axei*)
- Hookworm (*Bunostomum phlebotomum*)
- Large Bowel Worm (*Oesophagostomum radiatum*)
- Whipworm (*Trichuris spp.*)
- Thin Necked Intestinal Worm (*Nematodirus spathiger*)
- Intestinal Threadworm (*Strongyloides papillosus*)
- Lungworm (*Dictyocaulus viviparus*)
- Liver Fluke (*Fasciola hepatica*)

✓ SUCKING AND BITING LICE

- Long Nosed Sucking Lice (*Linognathus vituli*)
- Short Nosed Sucking Lice (*Haematopinus eurysternus*)
- Biting Lice (*Bovicola bovis*)
- Little Sucking Lice (*Solenopotes capillatus*)

✓ BUFFALO FLY (*Haematobia irritans exigua*)

✓ CATTLE TICK (*Boophilus microplus*)

RESTRAINTS

DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption.

CONTRAINDICATIONS

This product is contraindicated for use in calves under 16 weeks of age or less than 50 kg body weight.

This product should not be used in dogs, horses or any other pets.

This product is not recommended for use in goats as safety and efficacy have not been evaluated.

DOSAGE AND ADMINISTRATION

Vetsense Triclamec Ultra dose rate is 1 mL/10 kg body weight. Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

Over 650 kg dose at rate of 1 mL / 10 kg body weight. A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

GENERAL DIRECTIONS

Vetsense Triclamec Ultra is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product. A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions. Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian. After use clean all equipment with cooking oil. Do not use water.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT: **DO NOT USE** less than 49 days before slaughter for human consumption. Beef calves born to cows that were treated with Vetsense Triclamec Ultra during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the cow.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): **DO NOT USE** less than 140 days before slaughter for export. This ESI was correct at the time of label approval. Before using this product confirm the current ESI from Vetsense (0245779911) or the APVMA website (www.apvma.gov.au/residues). Beef calves born to cows that were treated with Vetsense Triclamec Ultra during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the cow.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash goggles and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY

Additional information is listed in the Safety Data Sheet.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

For containers less than or equal to 1L:

Dispose of container by wrapping with paper and putting in garbage.

For large containers greater than 1L:

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If the container has the drumMUSTER logo visible, has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.


STORAGE

Store below 30°C (room temperature).

APVMA No.: 94653/143094

LIVELIGHT	DOSE VOLUME	1L PACK TREATS	2.5L PACK TREATS	5L PACK TREATS	10L PACK TREATS
51-75 kg	7.5 mL	133	333	666	1333
76-100 kg	10 mL	100	250	500	1000
101-125 kg	12.5 mL	80	200	400	800
126-150 kg	15 mL	66	166	333	666
151-175 kg	17.5 mL	57	142	285	570
176-200 kg	20 mL	50	125	250	500
201-250 kg	25 mL	40	100	200	400
251-300 kg	30 mL	33	83	166	332
301-350 kg	35 mL	28	71	142	284
351-400 kg	40 mL	25	62	125	250
401-450 kg	45 mL	22	55	111	222
451-500 kg	50 mL	20	50	100	200
501-550 kg	55 mL	18	45	90	180
551-600 kg	60 mL	16	41	83	166
601-650 kg	65 mL	15	38	76	152

1L-Pack Application Guide

STEP 1	STEP 2
<p>Turn bottle upside down to fill the dosing chamber.</p> 	<p>Once chamber is filled, stand bottle upright to level the dosage.</p> 
STEP 3	STEP 4
<p>Tilt bottle to adjust to the correct dosage.</p> 	<p>Once you have the correct dosage, remove the cap.</p> 
STEP 5	STEP 6
<p>Pour out the liquid.</p> 	<p>Apply cap to reseal the bottle.</p> 