



Product Name: Vetsense Triclamec Ultra Pour-On Roundworm, Liver Fluke and External Parasiticide for Cattle
APVMA Approval No: 94653/143094

Label Name:	Vetsense Triclamec Ultra Pour-On Roundworm, Liver Fluke and External Parasiticide for Cattle
-------------	--

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS
------------------	--

Constituent Statements:	300 mg/mL TRICLABENDAZOLE 5 mg/mL ABAMECTIN Solvent: 500 mg/mL Diethylene glycol monobutyl ether
-------------------------	---

Claims:	<p>For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.</p> <p>Vetsense Triclamec Ultra contains Abamectin, a member of the macrocyclic lactones family of chemicals. It is effective against sensitive strains of internal and external parasites. It also contains Triclabendazole, a member of the benzimidazoles family of chemicals, effective against liver fluke.</p> <p>Indications</p> <p>For the treatment and control of adult and immature stages of:</p> <p>Gastro-intestinal roundworms:</p> <ul style="list-style-type: none">Barber's Pole Worm (<i>Haemonchus placei</i>)Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages)Small Intestinal Worm (<i>Cooperia oncophora</i>) (<i>Cooperia punctata</i>)Stomach Hair Worm (<i>Trichostrongylus axei</i>)Hookworm (<i>Bunostomum phlebotomum</i>)Large Bowel Worm (<i>Oesophagostomum radiatum</i>)Whipworm (<i>Trichuris spp.</i>)Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>)Intestinal Threadworm (<i>Strongyloides papillosus</i>)Lungworm (<i>Dictyocaulus viviparus</i>)Liver Fluke (<i>Fasciola hepatica</i>) including adult (12 week old), immature (6 week old) and early immature (4 week old) larvae. <p>Sucking and Biting Lice</p> <p>Long Nosed Sucking Lice (<i>Linognathus vituli</i>)</p>
---------	--

Short Nosed Sucking Lice (*Haematopinus eurysternus*)
 Biting Lice (*Bovicola bovis*)
 Little Sucking Lice (*Solenopotes capillatus*)
 Aids in control of:
 Buffalo Fly (*Haematobia irritans exigua*)
 Cattle Tick (*Boophilus microplus*)
 (Not recommended in strategic dipping programs for cattle tick control)
 Resistance may develop to any chemical.

Net Contents:	1 L 2 L 2.5 L 3 L 5 L 5.5 L 10 L 15 L
---------------	--

Directions for Use:	
---------------------	--

Restraints:	DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption.
-------------	---

Contraindications:	This product is contraindicated for use in calves under 16 weeks of age or less than 50 kg body weight. This product should not be used in dogs, horses or any other pets. This product is not recommended for use in goats as safety and efficacy have not been evaluated.
--------------------	---

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	Vetsense Triclamec Ultra dose rate is 1 mL / 10 kg body weight. Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected. [dosage chart] Over 650 kg dose at rate of 1 mL / 10 kg body weight. A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. This section contains file attachment.
----------------------------	--

General Directions:	<p>Vetsense Triclamec Ultra is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.</p> <p>A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. Do not brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions.</p> <p>Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian.</p> <p>After use clean all equipment with cooking oil. Do not use water.</p> <p>NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.</p>
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 49 days before slaughter for human consumption. Beef calves born to cows that were treated with Vetsense Triclamec Ultra during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the cow.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI):</p> <p>DO NOT use less than 140 days before slaughter for export. This ESI was correct at the time of label approval. Before using this product confirm the current ESI from Vetsense (02 4577 9911) or the APVMA website (www.apvma.gov.au/residues).</p> <p>Beef calves born to cows that were treated with Vetsense Triclamec Ultra during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the cow.</p>
---------------	---

Safety Directions:	<p>Harmful if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash goggles and contaminated clothing.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>Additional information is listed in the Safety Data Sheet.</p>
-------------------------	---

Environmental Statements:	<p>ENVIRONMENTAL PROTECTION Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
---------------------------	---

Disposal:	<p>[1 L containers]</p>
-----------	-------------------------

Dispose of container by wrapping with paper and putting in garbage.

[Other pack sizes]

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage:

Store below 30 °C (Room Temperature).

Dosage and Administration:

Live weight (kg)	Dose Volume (mL)
51-75	7.5
76-100	10
101-125	12.5
126-150	15
151-175	17.5
176-200	20
201-250	25
251-300	30
301-350	35
351-400	40
401-450	45
451-500	50
501-550	55
551-600	60
601-650	65