



Product Name: VETSENSE Abamec LV Pour-On for Cattle
APVMA Approval No: 95453/145718

Label Name:	VETSENSE Abamec LV Pour-On for Cattle
-------------	---------------------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	10 g/L ABAMECTIN
-------------------------	------------------

Claims:	<p>FOR THE TREATMENT AND CONTROL OF ABAMECTIN SENSITIVE INTERNAL AND EXTERNAL PARASITES OF DAIRY AND BEEF CATTLE.</p> <p>VETSENSE Abamec LV Pour-On for Cattle: Contains Abamectin (10g/L), a member of the macrocyclic lactones. It is effective against abamectin sensitive strains of internal and external parasites when applied to the skin along the backline of cattle. Resistance may develop to any chemical.</p> <p>Indications: For the treatment and control of adult and immature stages of: Gastro-intestinal roundworms: <i>Haemonchus placei</i> Barber's pole worm <i>Ostertagia ostertagi</i> (including inhibited stages) Small brown stomach worm <i>Cooperia oncophora</i>, <i>Cooperia punctata</i> Small intestinal worm <i>Trichostrongylus axei</i> Stomach hair worm <i>Bunostomum phlebotomum</i> Hookworm <i>Oesophagostomum radiatum</i> Nodule worm <i>Trichuris spp</i> Whipworm <i>Nematodirus spathiger</i> Thin necked intestinal worm <i>Strongyloides papillosum</i> Intestinal threadworm <i>Dictyocaulus viviparus</i> Large lungworm</p> <p>Sucking and Biting Lice: <i>Linognathus vituli</i> Long nosed sucking lice <i>Haematopinus eurysternus</i> Short nosed sucking lice <i>Bovicola bovis</i> Biting lice <i>Solenopotes capillatus</i> Little sucking lice <i>Boophilus microplus</i></p> <p>Cattle tick: aids in control (Not recommended in strategic dipping programs for cattle tick control).</p> <p>Sustained Activity:</p> <p>At the recommended dose rate, VETSENSE Abamec LV Pour-On for Cattle effectively controls infection with <i>Ostertagia spp.</i> (Small brown stomach worm) acquired up to 14 days after treatment, with <i>Dictyocaulus viviparus</i> (Lungworm) acquired up to 28 days after treatment.</p>
---------	---

Net Contents:	1L 2L 2.5L 5L 8L 10L
---------------	-------------------------------------

Directions for Use:	
---------------------	--

Restraints:	
-------------	--

Contraindications:	This product is contraindicated for use in calves under 16 weeks of age or less than 50 kg body weight This product should not be used in dogs, horses or any other pets This product is not recommended for use in goats as safety and efficacy have not been evaluated
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	Apply to the skin along the backline of the cattle at the dose rate of 1mL of VETSENSE Abamec LV Pour-On for Cattle/20kg bodyweight. Bodyweight Dose 51-100 kg 5.0 mL 101-150 kg 7.5 mL 151-200 kg 10.0 mL 201-250 kg 12.5 mL 251-300 kg 15.0 mL 301-350 kg 17.5 mL 351-400 kg 20.0 mL 401-450 kg 22.5 mL 451-500 kg 25.0 mL 501-550 kg 27.5 mL 551-600 kg 30.0 mL 601-650 kg 32.5 mL Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into 2 or more lines based on body weight, to avoid excessive overdosing.
----------------------------	---

General Directions:	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION
---------------------	--

Withholding Periods:	MEAT: DO NOT USE less than 35 days before slaughter for human consumption. MILK: Zero (0) Days
----------------------	---

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): An ESI has not been established for this product. Note: observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Vetsense on 0245779911 before using this product.
---------------	--

Safety Directions:	Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	Additional information is listed in the safety data sheet (SDS)
-------------------------	---

Environmental Statements:	ENVIRONMENTAL PROTECTION Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical and used containers.
---------------------------	--

Disposal:	For containers equal to 1L: Dispose of container by wrapping with paper and putting in garbage. For large containers greater than 1L: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------	--

Storage:	Store below 30 ° C (room temperature) in the closed, original container, in a dry, well ventilated area, and protect from sunlight.
----------	---