



Product Name: Trident Triple Combination Drench for Sheep
 APVMA Approval No: 90020/133591

Label Name:	Trident Triple Combination Drench for Sheep
-------------	---

Signal Headings:	<p>CAUTION</p> <p>KEEP OUT OF REACH OF CHILDREN</p> <p>FOR ANIMAL TREATMENT ONLY</p> <p>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
------------------	--

Constituent Statements:	<p>40 g/L LEVAMISOLE HYDROCHLORIDE</p> <p>1 g/L MOXIDECTIN</p> <p>25 g/L ALBENDAZOLE</p>
-------------------------	--

Claims:	<p>For the treatment and control of sensitive strains of gastrointestinal nematodes (roundworms), including nematodes that have single, dual or triple resistance to macrocyclic lactones (including abamectin resistant strains), benzimidazoles, imidazothiazoles (levamisole/morantel) or monepantel.</p> <p>TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP is a triple combination drench containing moxidectin, levamisole and albendazole, members respectively of the macrocyclic lactone ("mectin" drench), imidazothiazole (clear drench) and benzimidazole (white drench) chemical families.</p> <p>TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP provides treatment and control of sensitive and single, dual or triple resistant strains of the following gastrointestinal parasites of sheep:</p> <p>Barber's pole worm <i>Haemonchus contortus</i>*</p> <p>Stomach hair worm <i>Trichostrongylus axei</i>*</p> <p>Black scour worm <i>Trichostrongylus colubriformis</i>*, <i>Trichostrongylus rugatus</i>, <i>Trichostrongylus vitrinus</i></p> <p>Small brown stomach worm <i>Teladorsagia (Ostertagia) circumcincta</i>*, <i>Teladorsagia (Ostertagia) trifurcata</i>, <i>Teladorsagia (Ostertagia) davtiani</i></p> <p>Small intestinal worm <i>Copperia punctata</i>*, <i>Cooperia</i> spp.</p> <p>Thin necked intestinal worm <i>Nematodirus spathiger</i>, <i>Nematodirus</i> spp.</p> <p>Nodule worm <i>Oesophagostomum columbianum</i>, <i>Oesophagostomum</i> spp. Whipworm <i>Trichuris ovis</i></p> <p>*including immature fourth stage (L4) larvae</p> <p>Persistent Activity: TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP when used as recommended prevents reinfestation with ML-sensitive strains of Barber's pole worm (<i>Haemonchus contortus</i>) for 14 days and small brown stomach worm (<i>Teladorsagia (Ostertagia) circumcincta</i>) for 14 days.</p> <p>RESISTANCE AND WORM CONTROL PROGRAMS Resistance may develop to any chemical. However studies have shown that the use of combination drenches such as TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP can significantly delay the onset of resistance. Ask your local veterinarian or animal health adviser for recommended parasite</p>
---------	--

	management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.
--	---

Net Contents:	1 L 10 L 10.5 L 5 L
---------------	------------------------------

Directions for Use:	
---------------------	--

Restrains:	Do not re-treat animals for 17 days after last treatment. DO NOT USE in sheep which are producing or may in the future produce milk that may be used or processed for human consumption.
------------	---

Contraindications:	
--------------------	--

Precautions:	Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight, and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded, except under veterinary supervision. Doses of three or more times the recommended dose can cause symptoms of levamisole toxicity; hence bodyweights and doses should be determined accurately. Dehydrated animals may be more susceptible to toxicity.
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>SHAKE VIGOROUSLY FOR 30 SECONDS BEFORE USE</p> <p>For use as a quarantine drench: Treat all animals on arrival, or ideally at farm of origin, and hold in yards for 24 hours before release.</p> <p>For use in a strategic worm control program: Speak to your animal health adviser or veterinarian. TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP is ready to use. Administer orally 1 mL per 5 kg bodyweight using any standard drenching equipment. Check dose rates and equipment before drenching commences.</p> <p>DOSE RATE 1 mL/5 kg bodyweight to give a dose of 0.2 mg moxidectin, 6.8 mg levamisole and 5 mg albendazole per kg bodyweight.</p> <p>Body Weight (kg) Dose Volume (mL)</p> <table><tr><td>12 - 15</td><td>3</td></tr><tr><td>16 - 20</td><td>4</td></tr><tr><td>21 - 25</td><td>5</td></tr><tr><td>26 - 30</td><td>6</td></tr><tr><td>31 - 35</td><td>7</td></tr><tr><td>36 - 40</td><td>8</td></tr><tr><td>41 - 45</td><td>9</td></tr><tr><td>46 - 50</td><td>10</td></tr><tr><td>51 - 55</td><td>11</td></tr></table>	12 - 15	3	16 - 20	4	21 - 25	5	26 - 30	6	31 - 35	7	36 - 40	8	41 - 45	9	46 - 50	10	51 - 55	11
12 - 15	3																		
16 - 20	4																		
21 - 25	5																		
26 - 30	6																		
31 - 35	7																		
36 - 40	8																		
41 - 45	9																		
46 - 50	10																		
51 - 55	11																		

	<p>56 - 60 12 61 - 65 13 66 - 70 14 71 - 75 15</p> <p>For animals above 75 kg, dose at 1 mL per 5 kg bodyweight.</p> <p>A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by bodyweight in each group (ewes, rams, lambs). Do not underdose.</p> <p>Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.</p>
--	--

General Directions:	<p>TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP has a wide margin of safety when used as recommended and is readily accepted by sheep.</p> <p>TRIDENT TRIPLE COMBINATION DRENCH FOR SHEEP is safe to use concurrently with other treatments including mineral supplements, fluke treatments and vaccines.</p>
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 7 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE in sheep which are producing or may in the future produce milk that may be used or processed for human consumption.</p>
----------------------	--

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 17 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from either Jurox Pty Limited on 1800 023 312 or the APVMA website http://www.apvma.gov.au/residues.</p>
---------------	--

Safety Directions:	<p>Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear rubber gloves. Wash hands after use.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	<p>ENVIRONMENTAL PROTECTION</p> <p>Extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
---------------------------	---

Disposal:	<p>(1L) Dispose of container by wrapping with paper and putting in garbage.</p> <p>(5 L, 10 L, 10.5 L, 10 L + 5 L)</p>
-----------	--

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately. DO NOT burn empty containers or product.

Storage:

STORE BELOW 25°C (air conditioning). Protect from light.