

# TRIDENT<sup>®</sup>

40 g/L LEVAMISOLE HYDROCHLORIDE • 1 g/L MOXIDECTIN • 25 g/L ALBENDAZOLE

# NEW



- ▶ Triple combination sheep drench containing the potent active moxidectin
- ▶ Treatment and control of sensitive single, dual and triple resistant strains
- ▶ Broad Spectrum Parasite Control
- ▶ Short withholding periods for greater flexibility

## ONE DRENCH THREE-POINT DEFENCE



PROTECTION



CONTROL



CONFIDENCE

zoetis

# TRIDENT®

## TRIPLE COMBINATION DRENCH FOR SHEEP

Trident® Triple Combination Drench for Sheep is specially formulated for the effective treatment and control of sensitive strains of gastrointestinal roundworms, including those with single, dual, or triple resistance to:

- Macrocyclic lactones (including abamectin resistant strains)
- Benzimidazoles
- Imidazothiazoles (levamisole/morantel)
- Monepantel

### TRIPLE COMBINATION FORMULATION

Trident combines three proven active ingredients from different chemical families to maximize efficacy:

Active Ingredient	Chemical Family	Common Name
Moxidectin	Macrocyclic lactone	“Mectin” drench
Levamisole	Imidazothiazole	Clear drench
Albendazole	Benzimidazole	White drench

### BROAD SPECTRUM PARASITE CONTROL

Trident provides treatment and control of sensitive and resistant strains of the following gastrointestinal parasites of sheep, including immature larvae (L4):

- **Barber's pole worm** (*Haemonchus contortus*)
- **Stomach hair worm** (*Trichostrongylus axei*)
- **Black scour worm** (*Trichostrongylus colubriformis*, *T. rugatus*, *T. vitrinus*)
- **Small brown stomach worm** (*Teladorsagia circumcincta*, *T. trifurcata*, *T. davtiani*)
- **Small intestinal worm** (*Cooperia* spp., *Copperia punctata*)
- **Thin necked intestinal worm** (*Nematodirus* spp., *N. spathiger*)
- **Nodule worm** (*Oesophagostomum* spp., *O. columbianum*)
- **Whipworm** (*Trichuris ovis*)

### PERSISTENT ACTIVITY

When used as recommended, Trident prevents reinfestation with macrocyclic lactone-sensitive strains of:

- Barber's pole worm for **14 days**
- Small brown stomach worm for **14 days**

### DOSE RATE (1 mL/5 kg bodyweight)

Body Weight (kg)	Dose Volume (mL)
12 – 15	3
16 – 20	4
21 – 25	5
26 – 30	6
31 – 35	7
36 – 40	8
41 – 45	9
46 – 50	10
51 – 55	11
56 – 60	12
61 – 65	13
66 – 70	14
71 – 75	15
Above 75	1 mL per 5 kg bodyweight

### Withholding Period / Advice

<b>Meat</b>	<b>DO NOT USE</b> less than 7 days before slaughter for human consumption.
<b>Milk</b>	<b>DO NOT USE</b> in sheep producing or that may produce milk for human consumption or processing.
<b>Export Slaughter Interval (ESI)</b>	<b>DO NOT USE</b> less than 17 days before slaughter for export.