

NEW!

Control Barber's Pole
worm with a powerful,
dual-active oral drench



ABATAKTM DUAL

COMBINATION DRENCH FOR SHEEP

ACTIVE CONSTITUENTS: 50g/L CLOSANTEL, 1g/L ABAMECTIN

ABATAK™ DUAL

COMBINATION DRENCH FOR SHEEP

ACTIVE CONSTITUENTS: 50g/L CLOSANTEL, 1g/L ABAMECTIN

NEW!

For the control and treatment of roundworms, nasal bot, itch mite and mature and late immature liver fluke in sheep with activity against resistant strains of Barber's Pole Worm (including strains resistant to both Abamectin and Closantel) and 6-week sustained activity against Closantel susceptible Barber's Pole Worm (including strains resistant to macrocyclic lactones).

Pack sizes available

5L Backpack
10L Drum



About Barber's Pole worm*

Barber's Pole worm (*Haemonchus contortus*) is a blood-sucking roundworm that primarily affects sheep and goats. Here are some key points about it:

Lifecycle¹ - The worm's lifecycle involves eggs being passed in the sheep's feces, which hatch and develop into an infectious larval form.

Clinical signs² - Symptoms typically appear 3-6 weeks after ideal conditions (rain and warm weather) for the worms, which can lead to anemia and other health issues in sheep.


Appearance³ - The adult female worms have a characteristic "barber's pole" appearance due to their pink (blood-filled) intestinal tract twisted around the paler reproductive tract.

Ideal conditions⁴ - Barber's Pole worms thrive in moist and warm conditions, often seen in wet summers and autumn, making them a significant threat to young sheep.

Control Barber's Pole worm with a powerful, dual-active oral drench



- ✿ Broad Spectrum Drench against resistant strains of Barber's Pole Worm
- ✿ Dual Action parasite control
- ✿ Effective against internal parasites of sheep resistant to the single actives
- ✿ Broad-spectrum protection - Controls major roundworms, lungworms and liver fluke
- ✿ Prevents sheep re-infection for up to 6 weeks after drenching
- ✿ Limits pasture contamination with faecal eggs for up to 6 weeks after drenching
- ✿ Ewes may be treated at any stage of pregnancy
- ✿ Sustained activity against closantel-susceptible Barber's Pole Worms
- ✿ Administered orally using standard drenching equipment



	ABATAK™ DUAL	AVOMEK® DUEL
Active Constituents	50g/L CLOSANTEL, 1g/L ABAMECTIN	50g/L CLOSANTEL, 1g/L ABAMECTIN
Dose rate	1mL/5kg bodyweight	1mL/5kg bodyweight
Administration	Oral	Oral
Species	Sheep	Sheep
Withholding Periods		
Meat	49 days	49 days
Milk	Do not use in ewes, which are producing or may in the future produce milk or milk products for human consumption	Do not use in ewes, which are producing or may in the future produce milk or milk products for human consumption
ESI	84 days	84 days

ABATAK™ is a trademark of Abbey Laboratories Pty Ltd

AVOMEK® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH