



Australian Pesticides &
Veterinary Medicines Authority

Product Name: ABATAK DUAL COMBINATION DRENCH FOR SHEEP
APVMA Approval No: 95111/144720

Label Name:	ABATAK DUAL COMBINATION DRENCH FOR SHEEP
-------------	--

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	50 g/L closantel 1 g/L abamectin
-------------------------	-------------------------------------

Claims:	For the control & treatment of roundworms, nasal bot, itch mite and mature & late immature liver fluke in sheep with activity against resistant strains of Barber's Pole Worm (including strains resistant to both abamectin and closantel) and 6-week sustained activity against closantel susceptible Barber's Pole Worm (including strains resistant to macrocyclic lactones). ABATAK DUAL contains abamectin, a member of the macrocyclic lactone family of chemicals and Closantel, a member of the salicylanilide family of chemicals. INDICATIONS At the recommended dosage level, ABATAK DUAL provides treatment and control of the following sensitive strains of sheep parasites: Barber's Pole Worm (<i>Haemonchus contortus</i>) ABATAK DUAL is active against resistant strains of Barber's Pole Worm, including those resistant to BOTH closantel and abamectin. ABATAK DUAL also has sustained activity against closantel susceptible Barber's Pole Worm: • It prevents sheep re-infection for up to 6 weeks after drenching. • It also limits pasture contamination with faecal eggs for up to 6 weeks after drenching. Other adult and immature gastro-intestinal roundworms Small Brown Stomach Worm (<i>Ostertagia circumcincta</i>) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Black Scour Worm (<i>Trichostrongylus</i> spp.) Small Intestinal Worm (<i>Cooperia</i> spp.) Large Bowel Worm (<i>Oesophagostomum venulosum</i>)
---------	--

	Large Mouth Bowel Worm (<i>Chabertia ovina</i>) Nodule Worm (<i>Oesophagostomum columbianum</i>) Thin Necked Intestinal Worm (<i>Nematodirus</i> spp.) Whipworm (<i>Trichuris ovis</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>) Liver Fluke (<i>Fasciola hepatica</i>) mature and late immature (including 6 week old stages) Lungworm (<i>Dictyocaulus filaria</i>) Also for the control of Nasal Bot (<i>Oestrus ovis</i>) parasitic larval stages only Itch Mite (<i>Psorergates ovis</i>) Resistance may develop to any chemical. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench-resistance test be conducted before any drench is used.
--	---

Net Contents:	1 L 5 L 10 L 20 L
---------------	---

Directions for Use:	
---------------------	--

Restraints:	DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
-------------	---

Contraindications:	This product is contraindicated for use in lambs under 6 weeks of age or less than 10 kg body weight. This product should not be used in dogs, horses or any other pets. This product is not recommended for use in goats as safety and efficacy have not been evaluated.
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	SHAKE WELL BEFORE USE. Use the content within 3 months of first opening the container. Discard any unused portion. This product is administered orally using any standard drenching equipment that provides a consistent dose volume. Dose and equipment should be checked before drenching commences. The sheep dose is 1mL/5kg bodyweight (200µg Abamectin and 10mg Closantel per kg bodyweight). A representative sample of the animals should be weighed before treatment. Dose according to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two lines may be appropriate, to avoid excessive overdosing.
----------------------------	--

	This product is ready to use through standard equipment. Check accuracy of drenching equipment regularly. This section contains file attachment.
General Directions:	SAFETY TO SHEEP Ewes may be treated at any stage of pregnancy. Some sheep may cough slightly after treatment. This passing response is of no consequence. NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.
Withholding Periods:	MEAT: DO NOT USE less than 49 days before slaughter for human consumption.
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal health Pty Ltd on 020 8088 0720 or the APVMA website (www.apvma.gov.au/residues).
Safety Directions:	HARMFUL IF SWALLOWED. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If clothing becomes contaminated with product, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.
First Aid Instructions:	If poisoning occurs get to a doctor or hospital quickly.
First Aid Warnings:	
Additional User Safety:	CAN CAUSE PERMANENT VISION LOSS IF SWALLOWED. Additional information is in the Safety Data Sheet.
Environmental Statements:	ENVIRONMENTAL PROTECTION Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the product or used containers.
Disposal:	≤1L: Dispose of container by wrapping with paper and putting in garbage. >1L: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER collection site. If not recycling, break,

	crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
--	---

Storage:	Store below 30°C (room temperature). Protect from light.
----------	--

DOSE RATE TABLE:

Liveweight (Kg)	Dose Volume (mL)
11-15	3
16-20	4
21-25	5
26-30	6
31-35	7
36-40	8
41-50	10
51-60	12
61-70	14
71-80	16
Animals excess to 80kg to be dosed at 2mL/10kg bodyweight	