Australian Pesticides & Veterinary Medicines Authority

Company Name: DeltaX Pour-on for Cattle Lice and Fly Treatment
APVMA Approval No: 97518/153364

Label Name:	DeltaX Pour-on for Cattle Lice and Fly Treatment

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	15 g/L deltamethrin

Claims:	For control of Stable Fly (Stomoxys calcitrans), House Fly (Musca domestica) and Lice (Bovicola bovis, Linognathus vituli and Haematopinus eurysternus) and synthetic pyrethroid susceptible Buffalo Fly (Haematobia irritans exigua) on beef and dairy cattle. For control of synthetic pyrethroid susceptible Buffalo Fly (Haematobia irritans exigua) for up to 3 weeks and for control of Stable Fly (Stomoxys calcitrans) for up to 1 week on horses.

Net Contents:	1 L 2 L 5 L 6 L 10 L 20 L 25 L

Directions for Use:	

Restraints:	DO NOT USE on horses that may be used for human consumption.

Contraindications:	Contraindicated for use on animals that are severely stressed or in ill health. Not to be used on a horse's head forward of the ears. Not to be applied as a single spot.

Precautions:	Use with caution facial skin contact may cause temporary tingling/burning.

Side Effects:	

Dosage and Administration:	This section contains file attachment.

General Directions:	Cattle: To avoid operator insecticide contact when concurrent treatments are carried out, DeltaX Pour-On for Cattle Lice and Fly Treatment should be applied last. Operators must wear gloves. In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product. Apply with recommended applicator. The efficacy of this product was not affected by moderate rainfall when tested against cattle lice. Horses: Treatment with DeltaX will alleviate fly annoyance related behaviour including skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated with deltamethrin at 5 times the recommended dose rate. It is recommended that the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label. NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Withholding Periods:	CATTLE: MEAT: Zero (0) days MILK: Zero (0) days HORSES: NOT TO BE USED on horses that may be slaughtered for human consumption.

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website (www.apvma.gov.au/residues).

Safety Directions:	Product is poisonous if swallowed. May irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the product wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

First Aid Instructions:	If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:	

Additional User Safety:	Additional information is listed in the safety data sheet (SDS)

Environmental Statements:	DANGEROUS TO FISH. Do not contaminate dams, rivers or streams with the pesticide or used containers.

Disposal:	For Containers less than or equal to 1 L or 1 kg: Dispose of container by wrapping with paper and putting in garbage. For Containers greater than 1 L or 1 kg: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be recycled separately at the drumMUSTER collection site. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage:	Store below 30 °C (room temperature) in original container, tightly closed in a safe place.

DOSAGE AND ADMINISTRATION

Dose rate for Cattle:
5mL per 100kg liveweight.
The following table may be used as a guide

Pests Controlled	Rate (mL)	Live body weight	Critical comments
Cattle Lice (Bovicola bovis) (Linognathus vituli) (Haematopinus eurystemus)	5	Up to 100kg	Apply to animals in a race or drainage pen as a single strip along the backline. Retreat later in the season if reinfestation occurs.
	10	101 – 200kg	
	15	201 – 300kg	
Buffalo fly (Haematobia irritans exigua) Stable Fly (Stomoxys calcitrans) House fly (Musca domestica)	20	301 – 400kg	Apply to the whole herd- retreatment may be necessary
	25	401 – 500kg	
	30	501 – 600kg	
	For cattle over 600kg apply 5mL per 100kg liveweight		

Dose Rate for Horses:
5mL per 100kg liveweight.
The following table may be used as a guide.

Pests Controlled	Rate (mL)	Live body weight	Critical comments
Buffalo fly (Haematobia irritans exigua) Stable Fly (Stomoxys calcitrans)	5	Up to 100kg	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump. When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd — retreatment may be necessary
	10	101 – 200kg	
	15	201 – 300kg	
	20	301 – 400kg	
	25	401 – 500kg	
	30	501 – 600kg	
	For horses over 600kg apply 5mL per 100kg liveweight		